



PCT/IR 04 / 04 070
(09.12.04)



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

REC'D 09 DEC 2004
WIPO PCT


IB/04/4070

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed *P. Mahoney*
Dated *July 2004*

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

BEST AVAILABLE COPY



15DEC03 E859224-5 002093
P01/7700 0:00-0328907.1 ACCOUNT CHA


Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

12 DEC 2003



1. Your reference **PPD 70310/GB/P**

2. Patent application number
(The Patent Office will fill in this part) **0328907.1**

3. Full name, address and postcode of the or of each applicant (underline all surnames)
SYNGENTA PARTICIPATIONS AG
Intellectual Property Department
Schwarzwaldallee 215
4058 Basel
SWITZERLAND

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

8029555001

4. Title of the invention **CHEMICAL COMPOUNDS**

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

John Richard WATERMAN
Intellectual Property Department
Syngenta Limited
Jealott's Hill International Research Centre
PO Box 3538
Bracknell, Berkshire, RG42 6YA
UNITED KINGDOM

Patents ADP number (if you know it)

8029563001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

YES (b)

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form	81
Description	05
Claim(s)	01
Abstract	00
Drawing(s)	

SN

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.
SYNGENTA PARTICIPATIONS AG
Signature M A Rudd Date 12/12/03
Authorised Signatory

12. Name and daytime telephone number of person to contact in the United Kingdom

Margaret Ann RUDD 01344 413673

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes


- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

CHEMICAL COMPOUNDS

The present invention relates to spiroindoline derivatives, to processes for preparing them, to insecticidal, acaricidal, molluscicidal and nematocidal compositions comprising them and to methods of using them to combat and control insect, acarine, mollusc and nematode pests.

Spiroindoline derivatives with pharmaceutical properties are disclosed in for example US5763471, WO9825605, WO9429309, WO9828297 and WO9964002. Synthetic routes to selected compounds with pharmaceutical properties are described in Proc. Natl. Acad. Sci. USA (1995), 92, 7001, Tetrahedron (1997), 53, 10983 and Tetrahedron Letters (1997), 38, 1497. It has now surprisingly been found that certain spiroindolines have insecticidal properties.

The present invention therefore provides a method of combating and controlling insects, acarines, nematodes or molluscs which comprises applying to a pest, to a locus of a pest, or to a plant susceptible to attack by a pest an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a compound of formula (I):



wherein Y is a single bond, C=O, C=S or S(O)_m where m is 0, 1 or 2;

R¹ is hydrogen, optionally substituted alkyl, optionally substituted alkoxy carbonyl, optionally substituted alkyl carbonyl, aminocarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted aryl,

optionally substituted heteroaryl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted heterocycloxy, cyano, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, formyl, optionally substituted heterocyclyl, optionally substituted alkylthio, NO or $\text{NR}^{13}\text{R}^{14}$ where R^{13} and R^{14} are independently hydrogen, COR^{15} , optionally substituted alkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl or R^{13} and R^{14} together with the N atom to which they are attached form a group $-\text{N}=\text{C}(\text{R}^{16})-\text{NR}^{17}\text{R}^{18}$; R^{15} is H, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted aryl, optionally substituted aryloxy optionally substituted heteroaryl, optionally substituted heteroaryloxy or $\text{NR}^{19}\text{R}^{20}$; R^{16} , R^{17} and R^{18} are each independently H or lower alkyl; R^{19} and R^{20} are independently optionally substituted alkyl, optionally substituted aryl or optionally substituted heteroaryl;

R^2 and R^3 are independently hydrogen, halogen, cyano, optionally substituted alkyl, optionally substituted alkoxy or optionally substituted aryl;

each R^4 is independently halogen, nitro, cyano, optionally substituted C_{1-8} alkyl, optionally substituted C_{2-6} alkenyl, optionally substituted C_{2-6} alkynyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted C_{3-7} cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted alkylthio or $\text{R}^{21}\text{R}^{22}\text{N}$ where R^{21} and R^{22} are, independently, hydrogen, C_{1-8} alkyl, C_{3-7} cycloalkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-7} cycloalkyl(C_{1-4})alkyl, C_{2-6} haloalkyl, C_{1-6} alkoxy(C_{1-6})alkyl, C_{1-6} alkoxycarbonyl or R^{21} and R^{22} together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which may be optionally substituted by one or two C_{1-6} alkyl groups, or 2 adjacent groups R^4 together with the carbon atoms to which they are attached form a 4, 5, 6, or 7 membered carbocyclic or heterocyclic ring which may be optionally substituted by halogen; n is 0, 1, 2, 3 or 4;

Each R_a is independently hydrogen, halogen, hydroxy, cyano, optionally substituted C_{1-8} alkyl, optionally substituted C_{2-6} alkenyl, optionally substituted C_{2-6} alkynyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted C_{3-7} cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted alkylthio, optionally substituted arylthio or $R^{23}R^{24}N$ where R^{23} and R^{24} are, independently, hydrogen, C_{1-8} alkyl, C_{3-7} cycloalkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-7} cycloalkyl(C_{1-4})alkyl, C_{2-6} haloalkyl, C_{1-6} alkoxy(C_{1-6})alkyl, C_{1-6} alkoxycarbonyl or R^{23} and R^{24} together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which may be optionally substituted by one or two C_{1-6} alkyl groups, or two R_a groups attached to the same carbon atom are =O or two R_a groups attached to adjacent carbon atoms form a bond, or two R_a groups together with the carbon atom to which they are attached form a three- to seven-membered ring, that may be saturated or unsaturated, and that may contain one or two hetero atoms selected from the group consisting of N, O and S, and which may be optionally substituted by one or two C_{1-6} alkyl groups; or two R_a groups together form a group $-CH_2-$, $-CH=CH-$ or $-CH_2CH_2-$; p is 0, 1, 2, 3, 4, 5 or 6; q is 0, 1, 2, 3, 4, 5 or 6; provided that when p is 2 then q is not 2; p+q is 1, 2, 3, 4, 5 or 6;

R^8 is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted aryl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl or optionally substituted alkenylcarbonyl; or salts or N-oxides thereof.

The compounds of formula (I) may exist in different geometric or optical isomers or tautomeric forms. This invention covers all such isomers and tautomers and mixtures thereof in all proportions as well as isotopic forms such as deuterated compounds.

Each alkyl moiety either alone or as part of a larger group (such as alkoxy, alkoxycarbonyl, alkylcarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl) is a straight or

branched chain and is, for example, methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, isopropyl, n-butyl, sec-butyl, iso-butyl, tert-butyl or neo-pentyl. The alkyl groups are suitably C₁ to C₁₂ alkyl groups, but are preferably C₁-C₁₀, more preferably C₁-C₈, even more preferably preferably C₁-C₆ and most preferably C₁-C₄ alkyl groups.

- 5 When present, the optional substituents on an alkyl moiety (alone or as part of a larger group such as alkoxy, alkoxycarbonyl, alkylcarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl) include one or more of halogen, nitro, cyano, NCS-, C₃₋₇ cycloalkyl (itself optionally substituted with C₁₋₆ alkyl or halogen), C₅₋₇ cycloalkenyl (itself optionally substituted with C₁₋₆ alkyl or halogen), hydroxy, C₁₋₁₀ alkoxy, C₁₋₁₀ alkoxy(C₁₋₁₀)alkoxy, tri(C₁₋₄)alkylsilyl(C₁₋₆)alkoxy, C₁₋₆ alkoxycarbonyl(C₁₋₁₀)alkoxy, C₁₋₁₀ haloalkoxy, aryl(C₁₋₄)alkoxy (where the aryl group is optionally substituted), C₃₋₇ cycloalkyloxy (where the cycloalkyl group is optionally substituted with C₁₋₆ alkyl or halogen), C₂₋₁₀ alkenyloxy, C₂₋₁₀ alkynyloxy, SH, C₁₋₁₀ alkylthio, C₁₋₁₀ haloalkylthio, aryl(C₁₋₄)alkylthio (where the aryl group is optionally substituted), C₃₋₇ cycloalkylthio (where the cycloalkyl group is optionally substituted with C₁₋₆ alkyl or halogen), tri(C₁₋₄)alkylsilyl(C₁₋₆)alkylthio, arylthio (where the aryl group is optionally substituted), C₁₋₆ alkylsulfonyl, C₁₋₆ haloalkylsulfonyl, C₁₋₆ alkylsulfinyl, C₁₋₆ haloalkylsulfinyl, arylsulfonyl (where the aryl group may be optionally substituted), tri(C₁₋₄)alkylsilyl, aryl-di(C₁₋₄)alkylsilyl, (C₁₋₄)alkyldiarylsilyl, triarylsilyl, C₁₋₁₀ alkylcarbonyl, HO₂C, C₁₋₁₀ alkoxycarbonyl, aminocarbonyl, C₁₋₆ alkylaminocarbonyl, di(C₁₋₆ alkyl)aminocarbonyl, N-(C₁₋₃ alkyl)-N-(C₁₋₃ alkoxy)aminocarbonyl, C₁₋₆ alkylcarbonyloxy, arylcarbonyloxy (where the aryl group is optionally substituted), di(C₁₋₆)alkylaminocarbonyloxy, oximes such as =NOalkyl, =NOhaloalkyl and =NOaryl (itself optionally substituted), aryl (itself optionally substituted), heteroaryl (itself optionally substituted), heterocyclyl (itself optionally substituted with C₁₋₆ alkyl or halogen), aryloxy (where the aryl group is optionally substituted), heteroaryloxy, (where the heteroaryl group is optionally substituted), heterocyclyloxy (where the heterocyclyl group is optionally substituted with C₁₋₆ alkyl or halogen), amino, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, C₁₋₆ alkylcarbonylamino, N-(C₁₋₆)alkylcarbonyl-N-(C₁₋₆)alkylamino, C₂₋₆ alkenylcarbonyl, C₂₋₆ alkynylcarbonyl, C₃₋₆ alkenyloxycarbonyl, C₃₋₆ alkynyloxycarbonyl, aryloxycarbonyl (where

the aryl group is optionally substituted) and arylcarbonyl (where the aryl group is optionally substituted).

Alkenyl and alkynyl moieties can be in the form of straight or branched chains, and the alkenyl moieties, where appropriate, can be of either the (E)- or (Z)-configuration.

5 Examples are vinyl, allyl and propargyl.

When present, the optional substituents on alkenyl or alkynyl include those optional substituents given above for an alkyl moiety.

In the context of this specification acyl is optionally substituted C₁₋₆ alkylcarbonyl (for example acetyl), optionally substituted C₂₋₆ alkenylcarbonyl, optionally substituted C₂₋₆ alkynylcarbonyl, optionally substituted arylcarbonyl (for example benzoyl) or optionally substituted heteroarylcarbonyl.

Halogen is fluorine, chlorine, bromine or iodine.

Haloalkyl groups are alkyl groups which are substituted with one or more of the same or different halogen atoms and are, for example, CF₃, CF₂Cl, CF₃CH₂ or CHF₂CH₂.

15 In the context of the present specification the terms "aryl" and "aromatic ring system" refer to ring systems which may be mono-, bi- or tricyclic. Examples of such rings include phenyl, naphthalenyl, anthracenyl, indenyl or phenanthrenyl. A preferred aryl group is phenyl. In addition, the terms "heteroaryl", "heteroaromatic ring" or "heteroaromatic ring system" refer to an aromatic ring system containing at least one heteroatom and consisting
20 either of a single ring or of two or more fused rings. Preferably, single rings will contain up to three and bicyclic systems up to four heteroatoms which will preferably be chosen from nitrogen, oxygen and sulphur. Examples of such groups include furyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl,
25 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, 1,2,3-triazinyl, 1,2,4-triazinyl, 1,3,5-triazinyl, benzofuryl, benzisofuryl, benzothienyl, benzisothienyl, indolyl, isoindolyl, indazolyl, benzothiazolyl, benzisothiazolyl, benzoxazolyl, benzisoxazolyl, benzimidazolyl, 2,1,3-benzoxadiazole quinolinyl, isoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl,
30 quinoxalinyl, naphthyridinyl, benzotriazinyl, purinyl, pteridinyl and indolizinyl. Preferred

examples of heteroaromatic radicals include pyridyl, pyrimidyl, triazinyl, thienyl, furyl, oxazolyl, isoxazolyl, 2,1,3-benzoxadiazole and thiazolyl.

The terms heterocycle and heterocyclyl refer to a non-aromatic ring containing up to 10 atoms including one or more (preferably one or two) heteroatoms selected from O, S and

5 N. Examples of such rings include 1,3-dioxolane, tetrahydrofuran and morpholine.

When present, the optional substituents on heterocyclyl include C₁₋₆ alkyl and C₁₋₆ haloalkyl as well as those optional substituents given above for an alkyl moiety.

Cycloalkyl includes cyclopropyl, cyclopentyl and cyclohexyl.

Cycloalkenyl includes cyclopentenyl and cyclohexenyl.

10 When present, the optional substituents on cycloalkyl or cycloalkenyl include C₁₋₃ alkyl as well as those optional substituents given above for an alkyl moiety.

Carbocyclic rings include aryl, cycloalkyl and cycloalkenyl groups.

When present, the optional substituents on aryl or heteroaryl are selected independently, from halogen, nitro, cyano, NCS-, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy-
 15 (C₁₋₆)alkyl, C₂₋₆ alkenyl, C₂₋₆ haloalkenyl, C₂₋₆ alkynyl, C₃₋₇ cycloalkyl (itself optionally substituted with C₁₋₆ alkyl or halogen), C₅₋₇ cycloalkenyl (itself optionally substituted with C₁₋₆ alkyl or halogen), hydroxy, C₁₋₁₀ alkoxy, C₁₋₁₀ alkoxy(C₁₋₁₀)alkoxy, tri(C₁₋₄)alkyl-silyl(C₁₋₆)alkoxy, C₁₋₆ alkoxycarbonyl(C₁₋₁₀)alkoxy, C₁₋₁₀ haloalkoxy, aryl(C₁₋₄)alkoxy
 (where the aryl group is optionally substituted with halogen or C₁₋₆ alkyl), C₃₋₇ cycloalkyloxy
 20 (where the cycloalkyl group is optionally substituted with C₁₋₆ alkyl or halogen), C₂₋₁₀ alkenyloxy, C₂₋₁₀ alkynyloxy, SH, C₁₋₁₀ alkylthio, C₁₋₁₀ haloalkylthio, aryl(C₁₋₄)alkylthio C₃₋₇ cycloalkylthio (where the cycloalkyl group is optionally substituted with C₁₋₆ alkyl or halogen), tri(C₁₋₄)-alkylsilyl(C₁₋₆)alkylthio, arylthio, C₁₋₆ alkylsulfonyl, C₁₋₆ haloalkylsulfonyl, C₁₋₆ alkylsulfinyl, C₁₋₆ haloalkylsulfinyl, arylsulfonyl, tri(C₁₋₄)alkylsilyl,
 25 aryl-di(C₁₋₄)-alkylsilyl, (C₁₋₄)alkyl-diarylsilyl, triarylsilyl, C₁₋₁₀ alkylcarbonyl, HO₂C, C₁₋₁₀ alkoxycarbonyl, aminocarbonyl, C₁₋₆ alkylaminocarbonyl, di(C₁₋₆ alkyl)-aminocarbonyl, N-(C₁₋₃ alkyl)-N-(C₁₋₃ alkoxy)aminocarbonyl, C₁₋₆ alkylcarbonyloxy, arylcarbonyloxy, di(C₁₋₆)alkylamino-carbonyloxy, aryl (itself optionally substituted with C₁₋₆ alkyl or halogen), heteroaryl (itself optionally substituted with C₁₋₆ alkyl or halogen), heterocyclyl (itself
 30 optionally substituted with C₁₋₆ alkyl or halogen), aryloxy (where the aryl group is optionally

substituted with C₁₋₆ alkyl or halogen), heteroaryloxy (where the heteroaryl group is optionally substituted with C₁₋₆ alkyl or halogen), heterocyclyloxy (where the heterocyclyl group is optionally substituted with C₁₋₆ alkyl or halogen), amino, C₁₋₆ alkylamino, di(C₁₋₆)alkylamino, C₁₋₆ alkylcarbonylamino, N-(C₁₋₆)alkylcarbonyl-N-(C₁₋₆)alkylamino, arylcarbonyl, (where the aryl group is itself optionally substituted with halogen or C₁₋₆ alkyl) or two adjacent positions on an aryl or heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen or C₁₋₆ alkyl. Further substituents for aryl or heteroaryl include aryl carbonyl amino (where the aryl group is substituted by C₁₋₆ alkyl or halogen), (C₁₋₆)alkyloxycarbonylamino (C₁₋₆)alkyloxycarbonyl-N-(C₁₋₆)alkylamino, aryloxycarbonylamino (where the aryl group is substituted by C₁₋₆ alkyl or halogen), aryloxycarbonyl-N-(C₁₋₆)alkylamino, (where the aryl group is substituted by C₁₋₆ alkyl or halogen), arylsulphonylamino (where the aryl group is substituted by C₁₋₆ alkyl or halogen), arylsulphonyl-N-(C₁₋₆)alkylamino (where the aryl group is substituted by C₁₋₆ alkyl or halogen), aryl-N-(C₁₋₆)alkylamino (where the aryl group is substituted by C₁₋₆ alkyl or halogen), arylamino (where the aryl group is substituted by C₁₋₆ alkyl or halogen), heteroaryl amino (where the heteroaryl group is substituted by C₁₋₆ alkyl or halogen), heterocyclylamino (where the heterocyclyl group is substituted by C₁₋₆ alkyl or halogen), aminocarbonylamino, C₁₋₆ alkylaminocarbonyl amino, di(C₁₋₆)alkylaminocarbonyl amino, arylaminocarbonyl amino where the aryl group is substituted by C₁₋₆ alkyl or halogen), aryl-N-(C₁₋₆)alkylaminocarbonylamino where the aryl group is substituted by C₁₋₆ alkyl or halogen), C₁₋₆ alkylaminocarbonyl-N-(C₁₋₆)alkyl amino, di(C₁₋₆)alkylaminocarbonyl-N-(C₁₋₆)alkyl amino, arylaminocarbonyl-N-(C₁₋₆)alkyl amino where the aryl group is substituted by C₁₋₆ alkyl or halogen) and aryl-N-(C₁₋₆)alkylaminocarbonyl-N-(C₁₋₆)alkyl amino where the aryl group is substituted by C₁₋₆ alkyl or halogen).

For substituted phenyl moieties, heterocyclyl and heteroaryl groups it is preferred that one or more substituents are independently selected from halogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy(C₁₋₆)alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₁₋₆ alkylthio, C₁₋₆ haloalkylthio, C₁₋₆ alkylsulfinyl, C₁₋₆ haloalkylsulfinyl, C₁₋₆ alkylsulfonyl, C₁₋₆ haloalkylsulfonyl, C₂₋₆ alkenyl, C₂₋₆ haloalkenyl, C₂₋₆ alkynyl, C₃₋₇ cycloalkyl, nitro, cyano, CO₂H, C₁₋₆ alkylcarbonyl, C₁₋₆

alkoxycarbonyl, $R^{25}R^{26}N$ or $R^{27}R^{28}NC(O)$; wherein R^{25} , R^{26} , R^{27} and R^{28} are, independently, hydrogen or C_{1-6} alkyl. Further preferred substituents are aryl and heteroaryl groups.

Haloalkenyl groups are alkenyl groups which are substituted with one or more of the same or different halogen atoms.

5 It is to be understood that dialkylamino substituents include those where the dialkyl groups together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which is optionally substituted by one or two independently selected (C_{1-6})alkyl groups. When heterocyclic rings are formed by joining two groups on an N atom,
10 the resulting rings are suitably pyrrolidine, piperidine, thiomorpholine and morpholine each of which may be substituted by one or two independently selected (C_{1-6}) alkyl groups.

Preferably the optional substituents on an alkyl moiety include one or more of halogen, nitro, cyano, HO_2C , C_{1-10} alkoxy (itself optionally substituted by C_{1-10} alkoxy), aryl(C_{1-4})alkoxy, C_{1-10} alkylthio, C_{1-10} alkylcarbonyl, C_{1-10} alkoxycarbonyl, C_{1-6}
15 alkylaminocarbonyl, di(C_{1-6} alkyl)aminocarbonyl, (C_{1-6})alkylcarbonyloxy, optionally substituted phenyl, heteroaryl, aryloxy, arylcarbonyloxy, heteroaryloxy, heterocyclyl, heterocyclyloxy, C_{3-7} cycloalkyl (itself optionally substituted with (C_{1-6})alkyl or halogen), C_{3-7} cycloalkyloxy, C_{5-7} cycloalkenyl, C_{1-6} alkylsulfonyl, C_{1-6} alkylsulfinyl, tri(C_{1-4})alkylsilyl, tri(C_{1-4})alkylsilyl(C_{1-6})alkoxy, aryldi(C_{1-4})alkylsilyl, (C_{1-4})alkyldiarylsilyl and triarylsilyl.

20 Preferably the optional substituents on alkenyl or alkynyl include one or more of halogen, aryl and C_{3-7} cycloalkyl.

A preferred optional substituent for heterocyclyl is C_{1-6} alkyl.

Preferably the optional substituents for cycloalkyl include halogen, cyano and C_{1-3} alkyl.

25 Preferably the optional substituents for cycloalkenyl include C_{1-3} alkyl, halogen and cyano.

Preferably Y is a single bond, $C=O$ or $S(O)_m$ where m is 0, 1 or 2.

More preferably Y is a single bond, $C=O$ or SO_2 .

Yet more preferably Y is a single bond or $C=O$.

30 Most preferably Y is $C=O$.

Preferably R¹ is hydrogen, C₁₋₆ alkyl, C₁₋₆ cyanoalkyl, C₁₋₆ haloalkyl, C₃₋₇ cycloalkyl(C₁₋₄)alkyl, C₁₋₆ alkoxy(C₁₋₆)alkyl, heteroaryl(C₁₋₆)alkyl (wherein the heteroaryl group may be optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylthio, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), aryl(C₁₋₆)alkyl (wherein the aryl group may be optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylthio, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the aryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), C₁₋₆ alkylcarbonylamino(C₁₋₆)alkyl, aryl (which may be optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylthio, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), heteroaryl (which may be optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylthio, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, phenoxy (wherein the phenyl group is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), heteroaryloxy (optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy or C₁₋₆ haloalkoxy), heterocyclyloxy (optionally substituted by halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy or C₁₋₆ haloalkoxy), cyano, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₆ cycloalkyl, C₅₋₇ cycloalkenyl, heterocyclyl (optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy or C₁₋₆ haloalkoxy), C₁₋₆ alkylthio, C₁₋₆ haloalkylthio or NR¹³R¹⁴ where R¹³ and R¹⁴ are independently hydrogen, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy(C₁₋₆)alkyl, phenyl (which may be optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄

haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino, dialkylamino or C₁₋₄ alkoxy carbonyl), phenyl (C₁₋₆)alkyl (wherein the phenyl group may be optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino, dialkylamino, C₁₋₆ alkylsulfonyl, C₁₋₆ alkoxy carbonyl, or two adjacent positions on the phenyl ring may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), heteroaryl (C₁₋₆)alkyl (wherein the heteroaryl group may be optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylthio, C₁₋₆ alkoxy carbonyl, C₁₋₆ alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen) or heteroaryl (which may be optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy or C₁₋₆ haloalkoxy, C₁₋₄ alkoxy carbonyl C₁₋₆ alkylcarbonylamino, phenyloxy carbonylamino (wherein the phenyl group is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), amino, C₁₋₆ alkylamino or phenylamino (wherein the phenyl group is optionally substituted halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino)).

More preferably R¹ is C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy(C₁₋₆)alkyl, heteroaryl(C₁₋₃)alkyl (wherein the heteroaryl group may be optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₁₋₆ alkylsulfonyl, C₁₋₆ alkoxy carbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), phenyl(C₁₋₃)alkyl (wherein the phenyl group may be optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino, dialkylamino, C₁₋₆ alkylsulfonyl, C₁₋₆ alkoxy carbonyl, or two adjacent positions on the phenyl ring may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), phenyl (which may be optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino, dialkylamino, C₁₋₆ alkylsulfonyl, C₁₋₆ alkoxy carbonyl, or two adjacent positions on the phenyl ring may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic

ring, itself optionally substituted with halogen), heteroaryl (which may be optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₁₋₆ alkylsulfonyl, C₁₋₆ alkoxycarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, heterocyclyl (optionally substituted by halo, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy or C₁₋₆ haloalkoxy), C₁₋₆ alkylthio, C₁₋₆ haloalkylthio or NR¹³R¹⁴ where R¹³ and R¹⁴ are independently hydrogen, C₁₋₆ alkyl or C₁₋₆ haloalkyl, C₁₋₆ alkoxy(C₁₋₆)alkyl, C₂₋₆ alkylcarbonyl, phenylcarbonyl, (where the phenyl is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), phenyl(C₁₋₃)alkyl (wherein the phenyl group may be optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino, dialkylamino, C₁₋₆ alkylsulfonyl, C₁₋₆ alkoxycarbonyl, or two adjacent positions on the phenyl ring may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen) or heteroaryl(C₁₋₃)alkyl (wherein the heteroaryl group may be optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylthio, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen).

Even more preferably R¹ is C₁₋₆ alkyl, C₁₋₆ haloalkyl, heteroaryl(C₁₋₃)alkyl (wherein the heteroaryl group may be optionally substituted by halo, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl and where the heteroaryl group is a thiazole, pyridine, pyrimidine, pyrazine or pyridazine ring), heteroaryl (optionally substituted by halo, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl and where the heteroaryl group is a pyridine, pyrimidine, 2,1,3-benzoxadiazole, pyrazine or pyridazine ring), C₁₋₆ alkoxy, C₁₋₆ alkoxy(C₁₋₆)alkyl, C₁₋₆ alkylamino, C₁₋₆ alkoxy(C₁₋₆)alkylamino or heteroaryl(C₁₋₃)alkylamino (wherein the heteroaryl group may be optionally substituted by halo, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl and where the heteroaryl group is a thiazole, pyridine, pyrimidine, pyrazine or pyridazine ring).

Most preferably R^1 is pyridyl (optionally substituted by halo, C_{1-3} alkyl or C_{1-3} haloalkyl) especially halo-substituted pyridyl.

It is preferred that R^2 and R^3 are independently hydrogen, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy or cyano.

5 More preferably R^2 and R^3 are independently hydrogen, halogen, C_{1-2} alkyl, C_{1-2} haloalkyl, C_{1-2} alkoxy, cyano.

Even more preferably R^2 and R^3 are independently hydrogen or C_{1-4} alkyl.

Yet more preferably R^2 and R^3 are independently hydrogen or methyl.

Most preferably R^2 and R^3 are both hydrogen.

10 Preferably each R^4 is independently halogen, cyano, C_{1-8} alkyl, C_{1-8} haloalkyl, C_{1-6} cyanoalkyl, C_{1-6} alkoxy(C_{1-6})alkyl, C_{3-7} cycloalkyl(C_{1-6})alkyl, C_{5-6} cycloalkenyl(C_{1-6})alkyl, C_{3-6} alkenyloxy(C_{1-6})alkyl, C_{3-6} alkynyloxy(C_{1-6})alkyl, aryloxy(C_{1-6})alkyl, C_{1-6} carboxyalkyl, C_{1-6} alkylcarbonyl(C_{1-6})alkyl, C_{2-6} alkenylcarbonyl(C_{1-6})alkyl, C_{2-6} alkynylcarbonyl(C_{1-6})alkyl, C_{1-6} alkoxycarbonyl(C_{1-6})alkyl, C_{3-6} alkenyloxycarbonyl(C_{1-6})alkyl, C_{3-6} alkynyloxycarbonyl(C_{1-6})alkyl, aryloxycarbonyl(C_{1-6})alkyl, C_{1-6} alkylthio(C_{1-6})alkyl, C_{1-6} alkylsulfinyl(C_{1-6})alkyl, C_{1-6} alkylsulfonyl(C_{1-6})alkyl, aminocarbonyl(C_{1-6})alkyl, C_{1-6} alkylaminocarbonyl(C_{1-6})alkyl, di(C_{1-6})alkylaminocarbonyl(C_{1-6})alkyl, phenyl(C_{1-4})alkyl (wherein the phenyl group is optionally substituted by halogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkoxy, CN, NO_2 , aryl, heteroaryl, amino or dialkylamino),

20 heteroaryl(C_{1-4})alkyl (wherein the heteroaryl group is optionally substituted by halo, nitro, cyano, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy or C_{1-6} haloalkoxy), heterocyclyl(C_{1-4})alkyl (wherein the heterocyclyl group is optionally substituted by halo, nitro, cyano, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy or C_{1-6} haloalkoxy), C_{2-6} alkenyl, aminocarbonyl(C_{2-6})alkenyl, C_{1-6} alkylaminocarbonyl(C_{2-6})alkenyl, di(C_{1-6})alkylaminocarbonyl(C_{2-6})alkenyl, phenyl(C_{2-4})alkenyl, (wherein the phenyl group is optionally substituted by halogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkoxy, CN, NO_2 , aryl, heteroaryl, amino or dialkylamino), C_{2-6} alkynyl, trimethylsilyl(C_{2-6})alkynyl, aminocarbonyl(C_{2-6})alkynyl, C_{1-6} alkylaminocarbonyl(C_{2-6})alkynyl, di(C_{1-6})alkylaminocarbonyl(C_{2-6})alkynyl, C_{1-6} alkoxycarbonyl, C_{3-7} cycloalkyl, C_{3-7} halocycloalkyl, C_{3-7} cyanocycloalkyl, C_{1-3} alkyl(C_{3-7})cycloalkyl, C_{1-3} alkyl(C_{3-7})halocycloalkyl, phenyl (optionally substituted by halogen, C_{1-4}

30

alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), heteroaryl (optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy or C₁₋₆ haloalkoxy), heterocyclyl (wherein the heterocyclyl group is optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy or C₁₋₆ haloalkoxy), or 2 adjacent groups R⁴ together with the carbon atoms to which they are attached form a 4, 5, 6 or 7 membered carbocyclic or heterocyclic ring which may be optionally substituted by halogen, C₁₋₈ alkoxy, C₁₋₆ haloalkoxy, phenoxy (optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy or C₁₋₆ haloalkoxy), heteroaryloxy (optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy or C₁₋₆ haloalkoxy), C₁₋₈ alkylthio or R¹⁹R²⁰N where R¹⁹ and R²⁰ are, independently, hydrogen, C₁₋₈ alkyl, C₃₋₇ cycloalkyl, C₃₋₆ alkenyl, C₃₋₆ alkynyl, C₂₋₆ haloalkyl, C₁₋₆ alkoxycarbonyl or R¹⁹ and R²⁰ together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which may be optionally substituted by one or two C₁₋₆ alkyl groups; n is 0, 1, 2 or 3.

More preferably each R⁴ is independently halogen, cyano, C₁₋₈ alkyl, C₁₋₈ haloalkyl, C₁₋₈ cyanoalkyl, C₁₋₆ alkoxy(C₁₋₆)alkyl, C₂₋₆ alkynyl, trimethylsilyl(C₂₋₆)alkynyl, C₁₋₆ alkoxycarbonyl, C₃₋₇ cycloalkyl, C₁₋₃ alkyl (C₃₋₇) cycloalkyl, phenyl (optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), heterocyclyl (optionally substituted by halo, nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy or C₁₋₆ haloalkoxy), C₁₋₈ alkoxy, C₁₋₆ haloalkoxy, phenoxy (optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), heteroaryloxy (optionally substituted by halo, nitro, cyano, C₁₋₃ alkyl, C₁₋₃ haloalkyl, C₁₋₃ alkoxy or C₁₋₃ haloalkoxy), di(C₁₋₈)alkylamino, or 2 adjacent groups R⁴ together with the carbon atoms to which they are attached form a 4, 5, 6 or 7 membered carbocyclic or heterocyclic ring which may be optionally substituted by halogen; n is 0, 1, 2 or 3.

Even more preferably each R⁴ is independently halogen, cyano, C₁₋₈ alkyl, C₁₋₈ haloalkyl, C₁₋₈ cyanoalkyl, C₁₋₆ alkoxy(C₁₋₆)alkyl, C₂₋₆ alkynyl, heterocyclyl (optionally substituted by C₁₋₆ alkyl), C₁₋₈ alkoxy, C₁₋₆ haloalkoxy, phenoxy (optionally substituted by

halo, cyano, C₁₋₃ alkyl or C₁₋₃ haloalkyl), heteroaryloxy (optionally substituted by halo, cyano, C₁₋₃ alkyl or C₁₋₃ haloalkyl), di(C₁₋₈)alkylamino or 2 adjacent groups R⁴ together with the carbon atoms to which they are attached form a 4, 5, 6 or 7 membered carbocyclic or heterocyclic ring which may be optionally substituted by halogen; n is 0, 1, 2 or 3.

5 Yet more preferably each R⁴ is independently fluoro, chloro, bromo, cyano, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ cyanoalkyl or C₁₋₃ alkoxy(C₁₋₃)alkyl; n is 0, 1 or 2.

Most preferably each R⁴ is independently fluoro, chloro, bromo, C₁₋₄ alkyl or C₁₋₄ haloalkyl; n is 1 or 2.

Preferably R⁸ is C₁₋₁₀ alkyl, C₁₋₁₀ haloalkyl, aryl(C₁₋₆)alkyl (wherein the aryl group is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), heteroaryl(C₁₋₆)alkyl (wherein the heteroaryl group is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), arylcarbonyl-(C₁₋₆)alkyl (wherein the aryl group may be optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino and the alkyl group may be optionally substituted by aryl), C₂₋₈ alkenyl, C₂₋₈ haloalkenyl, aryl(C₂₋₆)-alkenyl (wherein the aryl group is optionally substituted halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino, C₁₋₆ alkoxy carbonyl, or two adjacent substituents can cyclise to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring), heteroaryl(C₂₋₆)-alkenyl (wherein the heteroaryl group is optionally substituted halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino, C₁₋₆ alkoxy carbonyl, or two adjacent substituents can cyclise to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring), C₂₋₆ alkynyl, phenyl(C₂₋₆)alkynyl (wherein the phenyl group is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), C₃₋₇ cycloalkyl, C₁₋₆ alkoxy carbonyl, C₁₋₆ alkyl carbonyl, C₁₋₆ haloalkyl carbonyl or aryl(C₂₋₆)alkenyl carbonyl (wherein the aryl group may be optionally substituted halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), or -C(R⁵¹)(R⁵²)-[CR⁵³=CR⁵⁴]_z-R⁵⁵ where z is 1 or 2, R⁵¹ and R⁵² are each independently H, halo or C₁₋₂ alkyl, R⁵³ and R⁵⁴ are each independently

30

H, halogen, C₁₋₄ alkyl or C₁₋₄ haloalkyl and R⁵⁵ is optionally substituted aryl or optionally substituted heteroaryl.

More preferably R⁸ is phenyl(C₁₋₄)alkyl (wherein the phenyl group is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), heteroaryl(C₁₋₆)alkyl (wherein the heteroaryl group is optionally substituted halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), phenyl(C₂₋₆)alkenyl (wherein the phenyl group is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino), heteroaryl(C₂₋₆)alkenyl (wherein the heteroaryl group is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino) or phenyl(C₂₋₆)alkynyl (wherein the phenyl group is optionally substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino, or – C(R⁵¹)(R⁵²)-[CR⁵³=CR⁵⁴]_z-R⁵⁵ where z is 1 or 2, R⁵¹ and R⁵² are each independently H, halo or C₁₋₂ alkyl, R⁵³ and R⁵⁴ are each independently H, halogen, C₁₋₄ alkyl or C₁₋₄ haloalkyl and R⁵⁵ is optionally substituted aryl or optionally substituted heteroaryl.

Most preferably R⁸ is –C(R⁵¹)(R⁵²)-[CR⁵³=CR⁵⁴]_z-R⁵⁵ where z is 1 or 2, preferably 1, R⁵¹ and R⁵² are each independently H, halo or C₁₋₂ alkyl, R⁵³ and R⁵⁴ are each independently H, halogen, C₁₋₄ alkyl or C₁₋₄ haloalkyl and R⁵⁵ is phenyl substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino or heteroaryl substituted by halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino.

R⁵¹ and R⁵² are preferably hydrogen.

R⁵³ and R⁵⁴ are preferably hydrogen or halogen, especially hydrogen.

R⁵⁵ is preferably phenyl substituted with one to three substituents selected from halogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, CN, NO₂, aryl, heteroaryl, amino or dialkylamino.

Preferably each R_a is independently hydrogen, halo, cyano, C₁₋₃ alkyl, hydroxy or two R_a groups together with the carbon atom to which they are attached form a carbonyl group

More preferably each R_a is independently hydrogen, fluoro, methyl, hydroxy or two R_a groups together with the carbon atom to which they are attached form a carbonyl group


Most preferably each R_a is hydrogen.

Preferably p is 1 or 2 and q is 2 or 3 and $p+q$ is 3, 4 or 5.

More preferably p is 1 or 2 and q is 2 or 3.

Most preferably p is 1 and q is 2 or 3.


Certain compounds of formula (I) are novel and as such form a further aspect of the invention. One group of novel compounds are compounds of formula I'



- 10 wherein Y is CO , R^2 and R^3 are both hydrogen and R^1 , R^4 , R^8 , R^a , n , p and q are as defined in relation to formula I provided that when n is 0, p is 1, q is 2, R^1 is CH_3 and all groups R_a are H then R^8 is not H, methyl, benzyl or $CH_2-CH=CH_2$ and when n is 0, $(CRa_2)_p$ is CH-phenyl, $(CRa_2)_q$ is $(CH_2)_2$ and R^1 is methyl then R^8 is not $COOCH_3$.

The compounds in Tables I to XLXIV below illustrate the compounds of the invention.

Table I provides 782 compounds of formula Ia



wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1

Table 1

Compound No	R ⁸	R ^{4a}	R ^{4b}	R ^{4c}	R ^{4d}
I-1	4-chlorobenzyl	H	H	H	H
I-2	Cinnamyl	H	H	H	H
I-3	4-chlorocinnamyl	H	H	H	H
I-4	4-fluorocinnamyl	H	H	H	H
I-5	4-bromocinnamyl	H	H	H	H
I-6	4-trifluoromethylcinnamyl	H	H	H	H
I-7	4-trifluoromethoxycinnamyl	H	H	H	H
I-8	4-pentafluoroethoxycinnamyl	H	H	H	H
I-9	4-methoxycinnamyl	H	H	H	H
I-10	4-ethoxycinnamyl	H	H	H	H
I-11	4-cyanocinnamyl	H	H	H	H
I-12	3-(6-chloro-pyridin-3-yl)-allyl	H	H	H	H
I-13	3-(4-chlorophenyl)-but-2-enyl	H	H	H	H
I-14	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	H	H
I-15	3-chloro-4-fluoro-cinnamyl	H	H	H	H
I-16	3,5-dichloro-cinnamyl	H	H	H	H
I-17	5-phenyl-penta-2,4-dienyl	H	H	H	H
I-18	4-isopropyloxycarbonylamino-cinnamyl	H	H	H	H
I-19	3-naphthalen-2-yl-allyl	H	H	H	H
I-20	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	H	H
I-21	3-(5-chloro-pyridin-2-yl)-allyl	H	H	H	H
I-22	3-pyridin-4-yl-allyl	H	H	H	H
I-23	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	H	H
I-24	4-chlorobenzyl	H	F	H	H
I-25	Cinnamyl	H	F	H	H
I-26	4-chlorocinnamyl	H	F	H	H

I-27	4-fluorocinnamyl	H	F	H	H
I-28	4-bromocinnamyl	H	F	H	H
I-29	4-trifluoromethylcinnamyl	H	F	H	H
I-30	4-trifluoromethoxycinnamyl	H	F	H	H
I-31	4-pentafluoroethoxycinnamyl	H	F	H	H
I-32	4-methoxycinnamyl	H	F	H	H
I-33	4-ethoxycinnamyl	H	F	H	H
I-34	4-cyanocinnamyl	H	F	H	H
I-35	3-(6-chloro-pyridin-3-yl)-allyl	H	F	H	H
I-36	3-(4-chlorophenyl)-but-2-enyl	H	F	H	H
I-37	3-(4-chlorophenyl)-3-fluoro-allyl	H	F	H	H
I-38	3-chloro-4-fluoro-cinnamyl	H	F	H	H
I-39	3,5-dichloro-cinnamyl	H	F	H	H
I-40	5-phenyl-penta-2,4-dienyl	H	F	H	H
I-41	4-isopropylloxycarbonylamino-cinnamyl	H	F	H	H
I-42	3-naphthalen-2-yl-allyl	H	F	H	H
I-43	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	F	H	H
I-44	3-(5-chloro-pyridin-2-yl)-allyl	H	F	H	H
I-45	3-pyridin-4-yl-allyl	H	F	H	H
I-46	3-(2-Chloro-pyridin-4-yl)-allyl	H	F	H	H
I-47	4-chlorobenzyl	H	Cl	H	H
I-48	Cinnamyl	H	Cl	H	H
I-49	4-chlorocinnamyl	H	Cl	H	H
I-50	4-fluorocinnamyl	H	Cl	H	H
I-51	4-bromocinnamyl	H	Cl	H	H
I-52	4-trifluoromethylcinnamyl	H	Cl	H	H
I-53	4-trifluoromethoxycinnamyl	H	Cl	H	H
I-54	4-pentafluoroethoxycinnamyl	H	Cl	H	H
I-55	4-methoxycinnamyl	H	Cl	H	H

I-56	4-ethoxycinnamyl	H	Cl	H	H
I-57	4-cyanocinnamyl	H	Cl	H	H
I-58	3-(6-chloro-pyridin-3-yl)-allyl	H	Cl	H	H
I-59	3-(4-chlorophenyl)-but-2-enyl	H	Cl	H	H
I-60	3-(4-chlorophenyl)-3-fluoro-allyl	H	Cl	H	H
I-61	3-chloro-4-fluoro-cinnamyl	H	Cl	H	H
I-62	3,5-dichloro-cinnamyl	H	Cl	H	H
I-63	5-phenyl-penta-2,4-dienyl	H	Cl	H	H
I-64	4-isopropoxyloxycarbonylamino-cinnamyl	H	Cl	H	H
I-65	3-naphthalen-2-yl-allyl	H	Cl	H	H
I-66	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	Cl	H	H
I-67	3-(5-chloro-pyridin-2-yl)-allyl	H	Cl	H	H
I-68	3-pyridin-4-yl-allyl	H	Cl	H	H
I-69	3-(2-Chloro-pyridin-4-yl)-allyl	H	Cl	H	H
I-70	4-chlorobenzyl	H	Br	H	H
I-71	Cinnamyl	H	Br	H	H
I-72	4-chlorocinnamyl	H	Br	H	H
I-73	4-fluorocinnamyl	H	Br	H	H
I-74	4-bromocinnamyl	H	Br	H	H
I-75	4-trifluoromethylcinnamyl	H	Br	H	H
I-76	4-trifluoromethoxycinnamyl	H	Br	H	H
I-77	4-pentafluoroethoxycinnamyl	H	Br	H	H
I-78	4-methoxycinnamyl	H	Br	H	H
I-79	4-ethoxycinnamyl	H	Br	H	H
I-80	4-cyanocinnamyl	H	Br	H	H
I-81	3-(6-chloro-pyridin-3-yl)-allyl	H	Br	H	H
I-82	3-(4-chlorophenyl)-but-2-enyl	H	Br	H	H
I-83	3-(4-chlorophenyl)-3-fluoro-allyl	H	Br	H	H
I-84	3-chloro-4-fluoro-cinnamyl	H	Br	H	H

I-85	3,5-dichloro-cinnamyl	H	Br	H	H
I-86	5-phenyl-penta-2,4-dienyl	H	Br	H	H
I-87	4-isopropoxyloxycarbonylamino-cinnamyl	H	Br	H	H
I-88	3-naphthalen-2-yl-allyl	H	Br	H	H
I-89	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	Br	H	H
I-90	3-(5-chloro-pyridin-2-yl)-allyl	H	Br	H	H
I-91	3-pyridin-4-yl-allyl	H	Br	H	H
I-92	3-(2-Chloro-pyridin-4-yl)-allyl	H	Br	H	H
I-93	4-chlorobenzyl	H	CN	H	H
I-94	Cinnamyl	H	CN	H	H
I-95	4-chlorocinnamyl	H	CN	H	H
I-96	4-fluorocinnamyl	H	CN	H	H
I-97	4-bromocinnamyl	H	CN	H	H
I-98	4-trifluoromethylcinnamyl	H	CN	H	H
I-99	4-trifluoromethoxycinnamyl	H	CN	H	H
I-100	4-pentafluoroethoxycinnamyl	H	CN	H	H
I-101	4-methoxycinnamyl	H	CN	H	H
I-102	4-ethoxycinnamyl	H	CN	H	H
I-103	4-cyanocinnamyl	H	CN	H	H
I-104	3-(6-chloro-pyridin-3-yl)-allyl	H	CN	H	H
I-105	3-(4-chlorophenyl)-but-2-enyl	H	CN	H	H
I-106	3-(4-chlorophenyl)-3-fluoro-allyl	H	CN	H	H
I-107	3-chloro-4-fluoro-cinnamyl	H	CN	H	H
I-108	3,5-dichloro-cinnamyl	H	CN	H	H
I-109	5-phenyl-penta-2,4-dienyl	H	CN	H	H
I-110	4-isopropoxyloxycarbonylamino-cinnamyl	H	CN	H	H
I-111	3-naphthalen-2-yl-allyl	H	CN	H	H
I-112	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	CN	H	H
I-113	3-(5-chloro-pyridin-2-yl)-allyl	H	CN	H	H

I-114	3-pyridin-4-yl-allyl	H	CN	H	H
I-115	3-(2-Chloro-pyridin-4-yl)-allyl	H	CN	H	H
I-116	4-chlorobenzyl	H	OMe	H	H
I-117	Cinnamyl	H	OMe	H	H
I-118	4-chlorocinnamyl	H	OMe	H	H
I-119	4-fluorocinnamyl	H	OMe	H	H
I-120	4-bromocinnamyl	H	OMe	H	H
I-121	4-trifluoromethylcinnamyl	H	OMe	H	H
I-122	4-trifluoromethoxycinnamyl	H	OMe	H	H
I-123	4-pentafluoroethoxycinnamyl	H	OMe	H	H
I-124	4-methoxycinnamyl	H	OMe	H	H
I-125	4-ethoxycinnamyl	H	OMe	H	H
I-126	4-cyanocinnamyl	H	OMe	H	H
I-127	3-(6-chloro-pyridin-3-yl)-allyl	H	OMe	H	H
I-128	3-(4-chlorophenyl)-but-2-enyl	H	OMe	H	H
I-129	3-(4-chlorophenyl)-3-fluoro-allyl	H	OMe	H	H
I-130	3-chloro-4-fluoro-cinnamyl	H	OMe	H	H
I-131	3,5-dichloro-cinnamyl	H	OMe	H	H
I-132	5-phenyl-penta-2,4-dienyl	H	OMe	H	H
I-133	4-isopropoxyloxycarbonylamino-cinnamyl	H	OMe	H	H
I-134	3-naphthalen-2-yl-allyl	H	OMe	H	H
I-135	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	OMe	H	H
I-136	3-(5-chloro-pyridin-2-yl)-allyl	H	OMe	H	H
I-137	3-pyridin-4-yl-allyl	H	OMe	H	H
I-138	3-(2-Chloro-pyridin-4-yl)-allyl	H	OMe	H	H
I-139	4-chlorobenzyl	H	OCF ₃	H	H
I-140	Cinnamyl	H	OCF ₃	H	H
I-141	4-chlorocinnamyl	H	OCF ₃	H	H
I-142	4-fluorocinnamyl	H	OCF ₃	H	H

I-143	4-bromocinnamyl	H	OCF ₃	H	H
I-144	4-trifluoromethylcinnamyl	H	OCF ₃	H	H
I-145	4-trifluoromethoxycinnamyl	H	OCF ₃	H	H
I-146	4-pentafluoroethoxycinnamyl	H	OCF ₃	H	H
I-147	4-methoxycinnamyl	H	OCF ₃	H	H
I-148	4-ethoxycinnamyl	H	OCF ₃	H	H
I-149	4-cyanocinnamyl	H	OCF ₃	H	H
I-150	3-(6-chloro-pyridin-3-yl)-allyl	H	OCF ₃	H	H
I-151	3-(4-chlorophenyl)-but-2-enyl	H	OCF ₃	H	H
I-152	3-(4-chlorophenyl)-3-fluoro-allyl	H	OCF ₃	H	H
I-153	3-chloro-4-fluoro-cinnamyl	H	OCF ₃	H	H
I-154	3,5-dichloro-cinnamyl	H	OCF ₃	H	H
I-155	5-phenyl-penta-2,4-dienyl	H	OCF ₃	H	H
I-156	4-isopropylloxycarbonylamino-cinnamyl	H	OCF ₃	H	H
I-157	3-naphthalen-2-yl-allyl	H	OCF ₃	H	H
I-158	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	OCF ₃	H	H
I-159	3-(5-chloro-pyridin-2-yl)-allyl	H	OCF ₃	H	H
I-160	3-pyridin-4-yl-allyl	H	OCF ₃	H	H
I-161	3-(2-Chloro-pyridin-4-yl)-allyl	H	OCF ₃	H	H
I-162	4-chlorobenzyl	H	CH ₃	H	H
I-163	Cinnamyl	H	CH ₃	H	H
I-164	4-chlorocinnamyl	H	CH ₃	H	H
I-165	4-fluorocinnamyl	H	CH ₃	H	H
I-166	4-bromocinnamyl	H	CH ₃	H	H
I-167	4-trifluoromethylcinnamyl	H	CH ₃	H	H
I-168	4-trifluoromethoxycinnamyl	H	CH ₃	H	H
I-169	4-pentafluoroethoxycinnamyl	H	CH ₃	H	H
I-170	4-methoxycinnamyl	H	CH ₃	H	H
I-171	4-ethoxycinnamyl	H	CH ₃	H	H

I-172	4-cyanocinnamyl	H	CH ₃	H	H
I-173	3-(6-chloro-pyridin-3-yl)-allyl	H	CH ₃	H	H
I-174	3-(4-chlorophenyl)-but-2-enyl	H	CH ₃	H	H
I-175	3-(4-chlorophenyl)-3-fluoro-allyl	H	CH ₃	H	H
I-176	3-chloro-4-fluoro-cinnamyl	H	CH ₃	H	H
I-177	3,5-dichloro-cinnamyl	H	CH ₃	H	H
I-178	5-phenyl-penta-2,4-dienyl	H	CH ₃	H	H
I-179	4-isopropoxyloxycarbonylamino-cinnamyl	H	CH ₃	H	H
I-180	3-naphthalen-2-yl-allyl	H	CH ₃	H	H
I-181	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	CH ₃	H	H
I-182	3-(5-chloro-pyridin-2-yl)-allyl	H	CH ₃	H	H
I-183	3-pyridin-4-yl-allyl	H	CH ₃	H	H
I-184	3-(2-Chloro-pyridin-4-yl)-allyl	H	CH ₃	H	H
I-185	4-chlorobenzyl	H	CF ₃	H	H
I-186	Cinnamyl	H	CF ₃	H	H
I-187	4-chlorocinnamyl	H	CF ₃	H	H
I-188	4-fluorocinnamyl	H	CF ₃	H	H
I-189	4-bromocinnamyl	H	CF ₃	H	H
I-190	4-trifluoromethylcinnamyl	H	CF ₃	H	H
I-191	4-trifluoromethoxycinnamyl	H	CF ₃	H	H
I-192	4-pentafluoroethoxycinnamyl	H	CF ₃	H	H
I-193	4-methoxycinnamyl	H	CF ₃	H	H
I-194	4-ethoxycinnamyl	H	CF ₃	H	H
I-195	4-cyanocinnamyl	H	CF ₃	H	H
I-196	3-(6-chloro-pyridin-3-yl)-allyl	H	CF ₃	H	H
I-197	3-(4-chlorophenyl)-but-2-enyl	H	CF ₃	H	H
I-198	3-(4-chlorophenyl)-3-fluoro-allyl	H	CF ₃	H	H
I-199	3-chloro-4-fluoro-cinnamyl	H	CF ₃	H	H
I-200	3,5-dichloro-cinnamyl	H	CF ₃	H	H

I-201	5-phenyl-penta-2,4-dienyl	H	CF ₃	H	H
I-202	4-isopropoxyloxycarbonylamino-cinnamyl	H	CF ₃	H	H
I-203	3-naphthalen-2-yl-allyl	H	CF ₃	H	H
I-204	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	CF ₃	H	H
I-205	3-(5-chloro-pyridin-2-yl)-allyl	H	CF ₃	H	H
I-206	3-pyridin-4-yl-allyl	H	CF ₃	H	H
I-207	3-(2-Chloro-pyridin-4-yl)-allyl	H	CF ₃	H	H
I-208	4-chlorobenzyl	H	H	Cl	H
I-209	Cinnamyl	H	H	Cl	H
I-210	4-chlorocinnamyl	H	H	Cl	H
I-211	4-fluorocinnamyl	H	H	Cl	H
I-212	4-bromocinnamyl	H	H	Cl	H
I-213	4-trifluoromethylcinnamyl	H	H	Cl	H
I-214	4-trifluoromethoxycinnamyl	H	H	Cl	H
I-215	4-pentafluoroethoxycinnamyl	H	H	Cl	H
I-216	4-methoxycinnamyl	H	H	Cl	H
I-217	4-ethoxycinnamyl	H	H	Cl	H
I-218	4-cyanocinnamyl	H	H	Cl	H
I-219	3-(6-chloro-pyridin-3-yl)-allyl	H	H	Cl	H
I-220	3-(4-chlorophenyl)-but-2-enyl	H	H	Cl	H
I-221	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	Cl	H
I-222	3-chloro-4-fluoro-cinnamyl	H	H	Cl	H
I-223	3,5-dichloro-cinnamyl	H	H	Cl	H
I-224	5-phenyl-penta-2,4-dienyl	H	H	Cl	H
I-225	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	Cl	H
I-226	3-naphthalen-2-yl-allyl	H	H	Cl	H
I-227	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	Cl	H
I-228	3-(5-chloro-pyridin-2-yl)-allyl	H	H	Cl	H
I-229	3-pyridin-4-yl-allyl	H	H	Cl	H

I-230	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	Cl	H
I-231	4-chlorobenzyl	H	H	F	H
I-232	Cinnamyl	H	H	F	H
I-233	4-chlorocinnamyl	H	H	F	H
I-234	4-fluorocinnamyl	H	H	F	H
I-235	4-bromocinnamyl	H	H	F	H
I-236	4-trifluoromethylcinnamyl	H	H	F	H
I-237	4-trifluoromethoxycinnamyl	H	H	F	H
I-238	4-pentafluoroethoxycinnamyl	H	H	F	H
I-239	4-methoxycinnamyl	H	H	F	H
I-240	4-ethoxycinnamyl	H	H	F	H
I-241	4-cyanocinnamyl	H	H	F	H
I-242	3-(6-chloro-pyridin-3-yl)-allyl	H	H	F	H
I-243	3-(4-chlorophenyl)-but-2-enyl	H	H	F	H
I-244	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	F	H
I-245	3-chloro-4-fluoro-cinnamyl	H	H	F	H
I-246	3,5-dichloro-cinnamyl	H	H	F	H
I-247	5-phenyl-penta-2,4-dienyl	H	H	F	H
I-248	4-isopropyloxycarbonylamino-cinnamyl	H	H	F	H
I-249	3-naphthalen-2-yl-allyl	H	H	F	H
I-250	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	F	H
I-251	3-(5-chloro-pyridin-2-yl)-allyl	H	H	F	H
I-252	3-pyridin-4-yl-allyl	H	H	F	H
I-253	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	F	H
I-254	4-chlorobenzyl	H	H	Br	H
I-255	Cinnamyl	H	H	Br	H
I-256	4-chlorocinnamyl	H	H	Br	H
I-257	4-fluorocinnamyl	H	H	Br	H
I-258	4-bromocinnamyl	H	H	Br	H

I-259	4-trifluoromethylcinnamyl	H	H	Br	H
I-260	4-trifluoromethoxycinnamyl	H	H	Br	H
I-261	4-pentafluoroethoxycinnamyl	H	H	Br	H
I-262	4-methoxycinnamyl	H	H	Br	H
I-263	4-ethoxycinnamyl	H	H	Br	H
I-264	4-cyanocinnamyl	H	H	Br	H
I-265	3-(6-chloro-pyridin-3-yl)-allyl	H	H	Br	H
I-266	3-(4-chlorophenyl)-but-2-enyl	H	H	Br	H
I-267	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	Br	H
I-268	3-chloro-4-fluoro-cinnamyl	H	H	Br	H
I-269	3,5-dichloro-cinnamyl	H	H	Br	H
I-270	5-phenyl-penta-2,4-dienyl	H	H	Br	H
I-271	4-isopropylloxycarbonylamino-cinnamyl	H	H	Br	H
I-272	3-naphthalen-2-yl-allyl	H	H	Br	H
I-273	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	Br	H
I-274	3-(5-chloro-pyridin-2-yl)-allyl	H	H	Br	H
I-275	3-pyridin-4-yl-allyl	H	H	Br	H
I-276	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	Br	H
I-277	4-chlorobenzyl	H	H	OCF ₃	H
I-278	Cinnamyl	H	H	OCF ₃	H
I-279	4-chlorocinnamyl	H	H	OCF ₃	H
I-280	4-fluorocinnamyl	H	H	OCF ₃	H
I-281	4-bromocinnamyl	H	H	OCF ₃	H
I-282	4-trifluoromethylcinnamyl	H	H	OCF ₃	H
I-283	4-trifluoromethoxycinnamyl	H	H	OCF ₃	H
I-284	4-pentafluoroethoxycinnamyl	H	H	OCF ₃	H
I-285	4-methoxycinnamyl	H	H	OCF ₃	H
I-286	4-ethoxycinnamyl	H	H	OCF ₃	H
I-287	4-cyanocinnamyl	H	H	OCF ₃	H

I-288	3-(6-chloro-pyridin-3-yl)-allyl	H	H	OCF ₃	H
I-289	3-(4-chlorophenyl)-but-2-enyl	H	H	OCF ₃	H
I-290	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	OCF ₃	H
I-291	3-chloro-4-fluoro-cinnamyl	H	H	OCF ₃	H
I-292	3,5-dichloro-cinnamyl	H	H	OCF ₃	H
I-293	5-phenyl-penta-2,4-dienyl	H	H	OCF ₃	H
I-294	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	OCF ₃	H
I-295	3-naphthalen-2-yl-allyl	H	H	OCF ₃	H
I-296	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	OCF ₃	H
I-297	3-(5-chloro-pyridin-2-yl)-allyl	H	H	OCF ₃	H
I-298	3-pyridin-4-yl-allyl	H	H	OCF ₃	H
I-299	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	OCF ₃	H
I-300	4-chlorobenzyl	H	H	CH ₃	H
I-301	Cinnamyl	H	H	CH ₃	H
I-302	4-chlorocinnamyl	H	H	CH ₃	H
I-303	4-fluorocinnamyl	H	H	CH ₃	H
I-304	4-bromocinnamyl	H	H	CH ₃	H
I-305	4-trifluoromethylcinnamyl	H	H	CH ₃	H
I-306	4-trifluoromethoxycinnamyl	H	H	CH ₃	H
I-307	4-pentafluoroethoxycinnamyl	H	H	CH ₃	H
I-308	4-methoxycinnamyl	H	H	CH ₃	H
I-309	4-ethoxycinnamyl	H	H	CH ₃	H
I-310	4-cyanocinnamyl	H	H	CH ₃	H
I-311	3-(6-chloro-pyridin-3-yl)-allyl	H	H	CH ₃	H
I-312	3-(4-chlorophenyl)-but-2-enyl	H	H	CH ₃	H
I-313	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	CH ₃	H
I-314	3-chloro-4-fluoro-cinnamyl	H	H	CH ₃	H
I-315	3,5-dichloro-cinnamyl	H	H	CH ₃	H
I-316	5-phenyl-penta-2,4-dienyl	H	H	CH ₃	H

I-317	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	CH ₃	H
I-318	3-naphthalen-2-yl-allyl	H	H	CH ₃	H
I-319	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	CH ₃	H
I-320	3-(5-chloro-pyridin-2-yl)-allyl	H	H	CH ₃	H
I-321	3-pyridin-4-yl-allyl	H	H	CH ₃	H
I-322	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	CH ₃	H
I-323	4-chlorobenzyl	H	H	CF ₃	H
I-324	Cinnamyl	H	H	CF ₃	H
I-325	4-chlorocinnamyl	H	H	CF ₃	H
I-326	4-fluorocinnamyl	H	H	CF ₃	H
I-327	4-bromocinnamyl	H	H	CF ₃	H
I-328	4-trifluoromethylcinnamyl	H	H	CF ₃	H
I-329	4-trifluoromethoxycinnamyl	H	H	CF ₃	H
I-330	4-pentafluoroethoxycinnamyl	H	H	CF ₃	H
I-331	4-methoxycinnamyl	H	H	CF ₃	H
I-332	4-ethoxycinnamyl	H	H	CF ₃	H
I-333	4-cyanocinnamyl	H	H	CF ₃	H
I-334	3-(6-chloro-pyridin-3-yl)-allyl	H	H	CF ₃	H
I-335	3-(4-chlorophenyl)-but-2-enyl	H	H	CF ₃	H
I-336	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	CF ₃	H
I-337	3-chloro-4-fluoro-cinnamyl	H	H	CF ₃	H
I-338	3,5-dichloro-cinnamyl	H	H	CF ₃	H
I-339	5-phenyl-penta-2,4-dienyl	H	H	CF ₃	H
I-340	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	CF ₃	H
I-341	3-naphthalen-2-yl-allyl	H	H	CF ₃	H
I-342	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	CF ₃	H
I-343	3-(5-chloro-pyridin-2-yl)-allyl	H	H	CF ₃	H
I-344	3-pyridin-4-yl-allyl	H	H	CF ₃	H
I-345	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	CF ₃	H

I-346	4-chlorobenzyl	F	H	H	H
I-347	Cinnamyl	F	H	H	H
I-348	4-chlorocinnamyl	F	H	H	H
I-349	4-fluorocinnamyl	F	H	H	H
I-350	4-bromocinnamyl	F	H	H	H
I-351	4-trifluoromethylcinnamyl	F	H	H	H
I-352	4-trifluoromethoxycinnamyl	F	H	H	H
I-353	4-pentafluoroethoxycinnamyl	F	H	H	H
I-354	4-methoxycinnamyl	F	H	H	H
I-355	4-ethoxycinnamyl	F	H	H	H
I-356	4-cyanocinnamyl	F	H	H	H
I-357	3-(6-chloro-pyridin-3-yl)-allyl	F	H	H	H
I-358	3-(4-chlorophenyl)-but-2-enyl	F	H	H	H
I-359	3-(4-chlorophenyl)-3-fluoro-allyl	F	H	H	H
I-360	3-chloro-4-fluoro-cinnamyl	F	H	H	H
I-361	3,5-dichloro-cinnamyl	F	H	H	H
I-362	5-phenyl-penta-2,4-dienyl	F	H	H	H
I-363	4-isopropoxyloxycarbonylamino-cinnamyl	F	H	H	H
I-364	3-naphthalen-2-yl-allyl	F	H	H	H
I-365	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	F	H	H	H
I-366	3-(5-chloro-pyridin-2-yl)-allyl	F	H	H	H
I-367	3-pyridin-4-yl-allyl	F	H	H	H
I-368	3-(2-Chloro-pyridin-4-yl)-allyl	F	H	H	H
I-369	4-chlorobenzyl	Cl	H	H	H
I-370	Cinnamyl	Cl	H	H	H
I-371	4-chlorocinnamyl	Cl	H	H	H
I-372	4-fluorocinnamyl	Cl	H	H	H
I-373	4-bromocinnamyl	Cl	H	H	H
I-374	4-trifluoromethylcinnamyl	Cl	H	H	H

I-375	4-trifluoromethoxycinnamyl	Cl	H	H	H
I-376	4-pentafluoroethoxycinnamyl	Cl	H	H	H
I-377	4-methoxycinnamyl	Cl	H	H	H
I-378	4-ethoxycinnamyl	Cl	H	H	H
I-379	4-cyanocinnamyl	Cl	H	H	H
I-380	3-(6-chloro-pyridin-3-yl)-allyl	Cl	H	H	H
I-381	3-(4-chlorophenyl)-but-2-enyl	Cl	H	H	H
I-382	3-(4-chlorophenyl)-3-fluoro-allyl	Cl	H	H	H
I-383	3-chloro-4-fluoro-cinnamyl	Cl	H	H	H
I-384	3,5-dichloro-cinnamyl	Cl	H	H	H
I-385	5-phenyl-penta-2,4-dienyl	Cl	H	H	H
I-386	4-isopropoxyloxycarbonylamino-cinnamyl	Cl	H	H	H
I-387	3-naphthalen-2-yl-allyl	Cl	H	H	H
I-388	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	Cl	H	H	H
I-389	3-(5-chloro-pyridin-2-yl)-allyl	Cl	H	H	H
I-390	3-pyridin-4-yl-allyl	Cl	H	H	H
I-391	3-(2-Chloro-pyridin-4-yl)-allyl	Cl	H	H	H
I-392	4-chlorobenzyl	Br	H	H	H
I-393	Cinnamyl	Br	H	H	H
I-394	4-chlorocinnamyl	Br	H	H	H
I-395	4-fluorocinnamyl	Br	H	H	H
I-396	4-bromocinnamyl	Br	H	H	H
I-397	4-trifluoromethylcinnamyl	Br	H	H	H
I-398	4-trifluoromethoxycinnamyl	Br	H	H	H
I-399	4-pentafluoroethoxycinnamyl	Br	H	H	H
I-400	4-methoxycinnamyl	Br	H	H	H
I-401	4-ethoxycinnamyl	Br	H	H	H
I-402	4-cyanocinnamyl	Br	H	H	H
I-403	3-(6-chloro-pyridin-3-yl)-allyl	Br	H	H	H

I-404	3-(4-chlorophenyl)-but-2-enyl	Br	H	H	H
I-405	3-(4-chlorophenyl)-3-fluoro-allyl	Br	H	H	H
I-406	3-chloro-4-fluoro-cinnamyl	Br	H	H	H
I-407	3,5-dichloro-cinnamyl	Br	H	H	H
I-408	5-phenyl-penta-2,4-dienyl	Br	H	H	H
I-409	4-isopropoxyloxycarbonylamino-cinnamyl	Br	H	H	H
I-410	3-naphthalen-2-yl-allyl	Br	H	H	H
I-411	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	Br	H	H	H
I-412	3-(5-chloro-pyridin-2-yl)-allyl	Br	H	H	H
I-413	3-pyridin-4-yl-allyl	Br	H	H	H
I-414	3-(2-Chloro-pyridin-4-yl)-allyl	Br	H	H	H
I-415	4-chlorobenzyl	CF ₃	H	H	H
I-416	Cinnamyl	CF ₃	H	H	H
I-417	4-chlorocinnamyl	CF ₃	H	H	H
I-418	4-fluorocinnamyl	CF ₃	H	H	H
I-419	4-bromocinnamyl	CF ₃	H	H	H
I-420	4-trifluoromethylcinnamyl	CF ₃	H	H	H
I-421	4-trifluoromethoxycinnamyl	CF ₃	H	H	H
I-422	4-pentafluoroethoxycinnamyl	CF ₃	H	H	H
I-423	4-methoxycinnamyl	CF ₃	H	H	H
I-424	4-ethoxycinnamyl	CF ₃	H	H	H
I-425	4-cyanocinnamyl	CF ₃	H	H	H
I-426	3-(6-chloro-pyridin-3-yl)-allyl	CF ₃	H	H	H
I-427	3-(4-chlorophenyl)-but-2-enyl	CF ₃	H	H	H
I-428	3-(4-chlorophenyl)-3-fluoro-allyl	CF ₃	H	H	H
I-429	3-chloro-4-fluoro-cinnamyl	CF ₃	H	H	H
I-430	3,5-dichloro-cinnamyl	CF ₃	H	H	H
I-431	5-phenyl-penta-2,4-dienyl	CF ₃	H	H	H
I-432	4-isopropoxyloxycarbonylamino-cinnamyl	CF ₃	H	H	H

I-433	3-naphthalen-2-yl-allyl	CF ₃	H	H	H
I-434	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	CF ₃	H	H	H
I-435	3-(5-chloro-pyridin-2-yl)-allyl	CF ₃	H	H	H
I-436	3-pyridin-4-yl-allyl	CF ₃	H	H	H
I-437	3-(2-Chloro-pyridin-4-yl)-allyl	CF ₃	H	H	H
I-438	4-chlorobenzyl	H	H	H	F
I-439	Cinnamyl	H	H	H	F
I-440	4-chlorocinnamyl	H	H	H	F
I-441	4-fluorocinnamyl	H	H	H	F
I-442	4-bromocinnamyl	H	H	H	F
I-443	4-trifluoromethylcinnamyl	H	H	H	F
I-444	4-trifluoromethoxycinnamyl	H	H	H	F
I-445	4-pentafluoroethoxycinnamyl	H	H	H	F
I-446	4-methoxycinnamyl	H	H	H	F
I-447	4-ethoxycinnamyl	H	H	H	F
I-448	4-cyanocinnamyl	H	H	H	F
I-449	3-(6-chloro-pyridin-3-yl)-allyl	H	H	H	F
I-450	3-(4-chlorophenyl)-but-2-enyl	H	H	H	F
I-451	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	H	F
I-452	3-chloro-4-fluoro-cinnamyl	H	H	H	F
I-453	3,5-dichloro-cinnamyl	H	H	H	F
I-454	5-phenyl-penta-2,4-dienyl	H	H	H	F
I-455	4-isopropylloxycarbonylamino-cinnamyl	H	H	H	F
I-456	3-naphthalen-2-yl-allyl	H	H	H	F
I-457	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	H	F
I-458	3-(5-chloro-pyridin-2-yl)-allyl	H	H	H	F
I-459	3-pyridin-4-yl-allyl	H	H	H	F
I-460	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	H	F
I-461	4-chlorobenzyl	H	H	H	Cl

I-462	Cinnamyl	H	H	H	Cl
I-463	4-chlorocinnamyl	H	H	H	Cl
I-464	4-fluorocinnamyl	H	H	H	Cl
I-465	4-bromocinnamyl	H	H	H	Cl
I-466	4-trifluoromethylcinnamyl	H	H	H	Cl
I-467	4-trifluoromethoxycinnamyl	H	H	H	Cl
I-468	4-pentafluoroethoxycinnamyl	H	H	H	Cl
I-469	4-methoxycinnamyl	H	H	H	Cl
I-470	4-ethoxycinnamyl	H	H	H	Cl
I-471	4-cyanocinnamyl	H	H	H	Cl
I-472	3-(6-chloro-pyridin-3-yl)-allyl	H	H	H	Cl
I-473	3-(4-chlorophenyl)-but-2-enyl	H	H	H	Cl
I-474	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	H	Cl
I-475	3-chloro-4-fluoro-cinnamyl	H	H	H	Cl
I-476	3,5-dichloro-cinnamyl	H	H	H	Cl
I-477	5-phenyl-penta-2,4-dienyl	H	H	H	Cl
I-478	4-isopropylloxycarbonylamino-cinnamyl	H	H	H	Cl
I-479	3-naphthalen-2-yl-allyl	H	H	H	Cl
I-480	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	H	Cl
I-481	3-(5-chloro-pyridin-2-yl)-allyl	H	H	H	Cl
I-482	3-pyridin-4-yl-allyl	H	H	H	Cl
I-483	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	H	Cl
I-484	4-chlorobenzyl	H	F	F	H
I-485	Cinnamyl	H	F	F	H
I-486	4-chlorocinnamyl	H	F	F	H
I-487	4-fluorocinnamyl	H	F	F	H
I-488	4-bromocinnamyl	H	F	F	H
I-489	4-trifluoromethylcinnamyl	H	F	F	H
I-490	4-trifluoromethoxycinnamyl	H	F	F	H

I-491	4-pentafluoroethoxycinnamyl	H	F	F	H
I-492	4-methoxycinnamyl	H	F	F	H
I-493	4-ethoxycinnamyl	H	F	F	H
I-494	4-cyanocinnamyl	H	F	F	H
I-495	3-(6-chloro-pyridin-3-yl)-allyl	H	F	F	H
I-496	3-(4-chlorophenyl)-but-2-enyl	H	F	F	H
I-497	3-(4-chlorophenyl)-3-fluoro-allyl	H	F	F	H
I-498	3-chloro-4-fluoro-cinnamyl	H	F	F	H
I-499	3,5-dichloro-cinnamyl	H	F	F	H
I-500	5-phenyl-penta-2,4-dienyl	H	F	F	H
I-501	4-isopropylloxycarbonylamino-cinnamyl	H	F	F	H
I-502	3-naphthalen-2-yl-allyl	H	F	F	H
I-503	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	F	F	H
I-504	3-(5-chloro-pyridin-2-yl)-allyl	H	F	F	H
I-505	3-pyridin-4-yl-allyl	H	F	F	H
I-506	3-(2-Chloro-pyridin-4-yl)-allyl	H	F	F	H
I-507	4-chlorobenzyl	H	F	Cl	H
I-508	Cinnamyl	H	F	Cl	H
I-509	4-chlorocinnamyl	H	F	Cl	H
I-510	4-fluorocinnamyl	H	F	Cl	H
I-511	4-bromocinnamyl	H	F	Cl	H
I-512	4-trifluoromethylcinnamyl	H	F	Cl	H
I-513	4-trifluoromethoxycinnamyl	H	F	Cl	H
I-514	4-pentafluoroethoxycinnamyl	H	F	Cl	H
I-515	4-methoxycinnamyl	H	F	Cl	H
I-516	4-ethoxycinnamyl	H	F	Cl	H
I-517	4-cyanocinnamyl	H	F	Cl	H
I-518	3-(6-chloro-pyridin-3-yl)-allyl	H	F	Cl	H
I-519	3-(4-chlorophenyl)-but-2-enyl	H	F	Cl	H

I-520	3-(4-chlorophenyl)-3-fluoro-allyl	H	F	Cl	H
I-521	3-chloro-4-fluoro-cinnamyl	H	F	Cl	H
I-522	3,5-dichloro-cinnamyl	H	F	Cl	H
I-523	5-phenyl-penta-2,4-dienyl	H	F	Cl	H
I-524	4-isopropoxyloxycarbonylamino-cinnamyl	H	F	Cl	H
I-525	3-naphthalen-2-yl-allyl	H	F	Cl	H
I-526	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	F	Cl	H
I-527	3-(5-chloro-pyridin-2-yl)-allyl	H	F	Cl	H
I-528	3-pyridin-4-yl-allyl	H	F	Cl	H
I-529	3-(2-Chloro-pyridin-4-yl)-allyl	H	F	Cl	H
I-530	4-chlorobenzyl	H	Cl	F	H
I-531	Cinnamyl	H	Cl	F	H
I-532	4-chlorocinnamyl	H	Cl	F	H
I-533	4-fluorocinnamyl	H	Cl	F	H
I-534	4-bromocinnamyl	H	Cl	F	H
I-535	4-trifluoromethylcinnamyl	H	Cl	F	H
I-536	4-trifluoromethoxycinnamyl	H	Cl	F	H
I-537	4-pentafluoroethoxycinnamyl	H	Cl	F	H
I-538	4-methoxycinnamyl	H	Cl	F	H
I-539	4-ethoxycinnamyl	H	Cl	F	H
I-540	4-cyanocinnamyl	H	Cl	F	H
I-541	3-(6-chloro-pyridin-3-yl)-allyl	H	Cl	F	H
I-542	3-(4-chlorophenyl)-but-2-enyl	H	Cl	F	H
I-543	3-(4-chlorophenyl)-3-fluoro-allyl	H	Cl	F	H
I-544	3-chloro-4-fluoro-cinnamyl	H	Cl	F	H
I-545	3,5-dichloro-cinnamyl	H	Cl	F	H
I-546	5-phenyl-penta-2,4-dienyl	H	Cl	F	H
I-547	4-isopropoxyloxycarbonylamino-cinnamyl	H	Cl	F	H
I-548	3-naphthalen-2-yl-allyl	H	Cl	F	H

I-549	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	Cl	F	H
I-550	3-(5-chloro-pyridin-2-yl)-allyl	H	Cl	F	H
I-551	3-pyridin-4-yl-allyl	H	Cl	F	H
I-552	3-(2-Chloro-pyridin-4-yl)-allyl	H	Cl	F	H
I-553	4-chlorobenzyl	H	Cl	Cl	H
I-554	Cinnamyl	H	Cl	Cl	H
I-555	4-chlorocinnamyl	H	Cl	Cl	H
I-556	4-fluorocinnamyl	H	Cl	Cl	H
I-557	4-bromocinnamyl	H	Cl	Cl	H
I-558	4-trifluoromethylcinnamyl	H	Cl	Cl	H
I-559	4-trifluoromethoxycinnamyl	H	Cl	Cl	H
I-560	4-pentafluoroethoxycinnamyl	H	Cl	Cl	H
I-561	4-methoxycinnamyl	H	Cl	Cl	H
I-562	4-ethoxycinnamyl	H	Cl	Cl	H
I-563	4-cyanocinnamyl	H	Cl	Cl	H
I-564	3-(6-chloro-pyridin-3-yl)-allyl	H	Cl	Cl	H
I-565	3-(4-chlorophenyl)-but-2-enyl	H	Cl	Cl	H
I-566	3-(4-chlorophenyl)-3-fluoro-allyl	H	Cl	Cl	H
I-567	3-chloro-4-fluoro-cinnamyl	H	Cl	Cl	H
I-568	3,5-dichloro-cinnamyl	H	Cl	Cl	H
I-569	5-phenyl-penta-2,4-dienyl	H	Cl	Cl	H
I-570	4-isopropylloxycarbonylamino-cinnamyl	H	Cl	Cl	H
I-571	3-naphthalen-2-yl-allyl	H	Cl	Cl	H
I-572	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	Cl	Cl	H
I-573	3-(5-chloro-pyridin-2-yl)-allyl	H	Cl	Cl	H
I-574	3-pyridin-4-yl-allyl	H	Cl	Cl	H
I-575	3-(2-Chloro-pyridin-4-yl)-allyl	H	Cl	Cl	H
I-576	4-chlorobenzyl	H	-OCF ₂ O-		H
I-577	Cinnamyl	H	-OCF ₂ O-		H

I-578	4-chlorocinnamyl	H	-OCF ₂ O-	H
I-579	4-fluorocinnamyl	H	-OCF ₂ O-	H
I-580	4-bromocinnamyl	H	-OCF ₂ O-	H
I-581	4-trifluoromethylcinnamyl	H	-OCF ₂ O-	H
I-582	4-trifluoromethoxycinnamyl	H	-OCF ₂ O-	H
I-583	4-pentafluoroethoxycinnamyl	H	-OCF ₂ O-	H
I-584	4-methoxycinnamyl	H	-OCF ₂ O-	H
I-585	4-ethoxycinnamyl	H	-OCF ₂ O-	H
I-586	4-cyanocinnamyl	H	-OCF ₂ O-	H
I-587	3-(6-chloro-pyridin-3-yl)-allyl	H	-OCF ₂ O-	H
I-588	3-(4-chlorophenyl)-but-2-enyl	H	-OCF ₂ O-	H
I-589	3-(4-chlorophenyl)-3-fluoro-allyl	H	-OCF ₂ O-	H
I-590	3-chloro-4-fluoro-cinnamyl	H	-OCF ₂ O-	H
I-591	3,5-dichloro-cinnamyl	H	-OCF ₂ O-	H
I-592	5-phenyl-penta-2,4-dienyl	H	-OCF ₂ O-	H
I-593	4-isopropoxyloxycarbonylamino-cinnamyl	H	-OCF ₂ O-	H
I-594	3-naphthalen-2-yl-allyl	H	-OCF ₂ O-	H
I-595	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	-OCF ₂ O-	H
I-596	3-(5-chloro-pyridin-2-yl)-allyl	H	-OCF ₂ O-	H
I-597	3-pyridin-4-yl-allyl	H	-OCF ₂ O-	H
I-598	3-(2-Chloro-pyridin-4-yl)-allyl	H	-OCF ₂ O-	H
I-599	4-chlorobenzyl	H	-C ₄ H ₄ -	H
I-600	Cinnamyl	H	-C ₄ H ₄ -	H
I-601	4-chlorocinnamyl	H	-C ₄ H ₄ -	H
I-602	4-fluorocinnamyl	H	-C ₄ H ₄ -	H
I-603	4-bromocinnamyl	H	-C ₄ H ₄ -	H
I-604	4-trifluoromethylcinnamyl	H	-C ₄ H ₄ -	H
I-605	4-trifluoromethoxycinnamyl	H	-C ₄ H ₄ -	H
I-606	4-pentafluoroethoxycinnamyl	H	-C ₄ H ₄ -	H

I-607	4-methoxycinnamyl	H	-C ₄ H ₄ -	H	
I-608	4-ethoxycinnamyl	H	-C ₄ H ₄ -	H	
I-609	4-cyanocinnamyl	H	-C ₄ H ₄ -	H	
I-610	3-(6-chloro-pyridin-3-yl)-allyl	H	-C ₄ H ₄ -	H	
I-611	3-(4-chlorophenyl)-but-2-enyl	H	-C ₄ H ₄ -	H	
I-612	3-(4-chlorophenyl)-3-fluoro-allyl	H	-C ₄ H ₄ -	H	
I-613	3-chloro-4-fluoro-cinnamyl	H	-C ₄ H ₄ -	H	
I-614	3,5-dichloro-cinnamyl	H	-C ₄ H ₄ -	H	
I-615	5-phenyl-penta-2,4-dienyl	H	-C ₄ H ₄ -	H	
I-616	4-isopropylloxycarbonylamino-cinnamyl	H	-C ₄ H ₄ -	H	
I-617	3-naphthalen-2-yl-allyl	H	-C ₄ H ₄ -	H	
I-618	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	-C ₄ H ₄ -	H	
I-619	3-(5-chloro-pyridin-2-yl)-allyl	H	-C ₄ H ₄ -	H	
I-620	3-pyridin-4-yl-allyl	H	-C ₄ H ₄ -	H	
I-621	3-(2-Chloro-pyridin-4-yl)-allyl	H	-C ₄ H ₄ -	H	
I-622	4-chlorobenzyl	Cl	H	Cl	H
I-623	Cinnamyl	Cl	H	Cl	H
I-624	4-chlorocinnamyl	Cl	H	Cl	H
I-625	4-fluorocinnamyl	Cl	H	Cl	H
I-226	4-bromocinnamyl	Cl	H	Cl	H
I-627	4-trifluoromethylcinnamyl	Cl	H	Cl	H
I-628	4-trifluoromethoxycinnamyl	Cl	H	Cl	H
I-629	4-pentafluoroethoxycinnamyl	Cl	H	Cl	H
I-630	4-methoxycinnamyl	Cl	H	Cl	H
I-631	4-ethoxycinnamyl	Cl	H	Cl	H
I-632	4-cyanocinnamyl	Cl	H	Cl	H
I-633	3-(6-chloro-pyridin-3-yl)-allyl	Cl	H	Cl	H
I-634	3-(4-chlorophenyl)-but-2-enyl	Cl	H	Cl	H
I-635	3-(4-chlorophenyl)-3-fluoro-allyl	Cl	H	Cl	H

I-636	3-chloro-4-fluoro-cinnamyl	Cl	H	Cl	H
I-637	3,5-dichloro-cinnamyl	Cl	H	Cl	H
I-638	5-phenyl-penta-2,4-dienyl	Cl	H	Cl	H
I-639	4-isopropylloxycarbonylamino-cinnamyl	Cl	H	Cl	H
I-640	3-naphthalen-2-yl-allyl	Cl	H	Cl	H
I-641	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	Cl	H	Cl	H
I-642	3-(5-chloro-pyridin-2-yl)-allyl	Cl	H	Cl	H
I-643	3-pyridin-4-yl-allyl	Cl	H	Cl	H
I-644	3-(2-Chloro-pyridin-4-yl)-allyl	Cl	H	Cl	H
I-645	4-chlorobenzyl	Cl	Cl	H	H
I-646	Cinnamyl	Cl	Cl	H	H
I-647	4-chlorocinnamyl	Cl	Cl	H	H
I-648	4-fluorocinnamyl	Cl	Cl	H	H
I-649	4-bromocinnamyl	Cl	Cl	H	H
I-650	4-trifluoromethylcinnamyl	Cl	Cl	H	H
I-651	4-trifluoromethoxycinnamyl	Cl	Cl	H	H
I-652	4-pentafluoroethoxycinnamyl	Cl	Cl	H	H
I-653	4-methoxycinnamyl	Cl	Cl	H	H
I-654	4-ethoxycinnamyl	Cl	Cl	H	H
I-655	4-cyanocinnamyl	Cl	Cl	H	H
I-656	3-(6-chloro-pyridin-3-yl)-allyl	Cl	Cl	H	H
I-657	3-(4-chlorophenyl)-but-2-enyl	Cl	Cl	H	H
I-658	3-(4-chlorophenyl)-3-fluoro-allyl	Cl	Cl	H	H
I-659	3-chloro-4-fluoro-cinnamyl	Cl	Cl	H	H
I-660	3,5-dichloro-cinnamyl	Cl	Cl	H	H
I-661	5-phenyl-penta-2,4-dienyl	Cl	Cl	H	H
I-662	4-isopropylloxycarbonylamino-cinnamyl	Cl	Cl	H	H
I-663	3-naphthalen-2-yl-allyl	Cl	Cl	H	H
I-664	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	Cl	Cl	H	H

I-665	3-(5-chloro-pyridin-2-yl)-allyl	Cl	Cl	H	H
I-666	3-pyridin-4-yl-allyl	Cl	Cl	H	H
I-667	3-(2-Chloro-pyridin-4-yl)-allyl	Cl	Cl	H	H
I-668	4-chlorobenzyl	H	Cl	H	Cl
I-669	Cinnamyl	H	Cl	H	Cl
I-670	4-chlorocinnamyl	H	Cl	H	Cl
I-671	4-fluorocinnamyl	H	Cl	H	Cl
I-672	4-bromocinnamyl	H	Cl	H	Cl
I-673	4-trifluoromethylcinnamyl	H	Cl	H	Cl
I-674	4-trifluoromethoxycinnamyl	H	Cl	H	Cl
I-675	4-pentafluoroethoxycinnamyl	H	Cl	H	Cl
I-676	4-methoxycinnamyl	H	Cl	H	Cl
I-677	4-ethoxycinnamyl	H	Cl	H	Cl
I-678	4-cyanocinnamyl	H	Cl	H	Cl
I-679	3-(6-chloro-pyridin-3-yl)-allyl	H	Cl	H	Cl
I-680	3-(4-chlorophenyl)-but-2-enyl	H	Cl	H	Cl
I-681	3-(4-chlorophenyl)-3-fluoro-allyl	H	Cl	H	Cl
I-682	3-chloro-4-fluoro-cinnamyl	H	Cl	H	Cl
I-683	3,5-dichloro-cinnamyl	H	Cl	H	Cl
I-684	5-phenyl-penta-2,4-dienyl	H	Cl	H	Cl
I-685	4-isopropylloxycarbonylamino-cinnamyl	H	Cl	H	Cl
I-686	3-naphthalen-2-yl-allyl	H	Cl	H	Cl
I-687	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	Cl	H	Cl
I-688	3-(5-chloro-pyridin-2-yl)-allyl	H	Cl	H	Cl
I-689	3-pyridin-4-yl-allyl	H	Cl	H	Cl
I-690	3-(2-Chloro-pyridin-4-yl)-allyl	H	Cl	H	Cl
I-691	4-chlorobenzyl	H	F	H	F
I-692	Cinnamyl	H	F	H	F
I-693	4-chlorocinnamyl	H	F	H	F


I-694	4-fluorocinnamyl	H	F	H	F
I-695	4-bromocinnamyl	H	F	H	F
I-696	4-trifluoromethylcinnamyl	H	F	H	F
I-697	4-trifluoromethoxycinnamyl	H	F	H	F
I-698	4-pentafluoroethoxycinnamyl	H	F	H	F
I-699	4-methoxycinnamyl	H	F	H	F
I-700	4-ethoxycinnamyl	H	F	H	F
I-701	4-cyanocinnamyl	H	F	H	F
I-702	3-(6-chloro-pyridin-3-yl)-allyl	H	F	H	F
I-703	3-(4-chlorophenyl)-but-2-enyl	H	F	H	F
I-704	3-(4-chlorophenyl)-3-fluoro-allyl	H	F	H	F
I-705	3-chloro-4-fluoro-cinnamyl	H	F	H	F
I-706	3,5-dichloro-cinnamyl	H	F	H	F
I-707	5-phenyl-penta-2,4-dienyl	H	F	H	F
I-708	4-isopropylloxycarbonylamino-cinnamyl	H	F	H	F
I-709	3-naphthalen-2-yl-allyl	H	F	H	F
I-710	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	F	H	F
I-711	3-(5-chloro-pyridin-2-yl)-allyl	H	F	H	F
I-712	3-pyridin-4-yl-allyl	H	F	H	F
I-713	3-(2-Chloro-pyridin-4-yl)-allyl	H	F	H	F
I-714	4-chlorobenzyl	F	H	F	H
I-715	Cinnamyl	F	H	F	H
I-716	4-chlorocinnamyl	F	H	F	H
I-717	4-fluorocinnamyl	F	H	F	H
I-718	4-bromocinnamyl	F	H	F	H
I-719	4-trifluoromethylcinnamyl	F	H	F	H
I-720	4-trifluoromethoxycinnamyl	F	H	F	H
I-721	4-pentafluoroethoxycinnamyl	F	H	F	H
I-722	4-methoxycinnamyl	F	H	F	H

I-723	4-ethoxycinnamyl	F	H	F	H
I-724	4-cyanocinnamyl	F	H	F	H
I-725	3-(6-chloro-pyridin-3-yl)-allyl	F	H	F	H
I-726	3-(4-chlorophenyl)-but-2-enyl	F	H	F	H
I-727	3-(4-chlorophenyl)-3-fluoro-allyl	F	H	F	H
I-728	3-chloro-4-fluoro-cinnamyl	F	H	F	H
I-729	3,5-dichloro-cinnamyl	F	H	F	H
I-730	5-phenyl-penta-2,4-dienyl	F	H	F	H
I-731	4-isopropylloxycarbonylamino-cinnamyl	F	H	F	H
I-732	3-naphthalen-2-yl-allyl	F	H	F	H
I-733	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	F	H	F	H
I-734	3-(5-chloro-pyridin-2-yl)-allyl	F	H	F	H
I-735	3-pyridin-4-yl-allyl	F	H	F	H
I-736	3-(2-Chloro-pyridin-4-yl)-allyl	F	H	F	H
I-737	4-chlorobenzyl	F	F	H	H
I-738	Cinnamyl	F	F	H	H
I-739	4-chlorocinnamyl	F	F	H	H
I-740	4-fluorocinnamyl	F	F	H	H
I-741	4-bromocinnamyl	F	F	H	H
I-742	4-trifluoromethylcinnamyl	F	F	H	H
I-743	4-trifluoromethoxycinnamyl	F	F	H	H
I-744	4-pentafluoroethoxycinnamyl	F	F	H	H
I-745	4-methoxycinnamyl	F	F	H	H
I-746	4-ethoxycinnamyl	F	F	H	H
I-747	4-cyanocinnamyl	F	F	H	H
I-748	3-(6-chloro-pyridin-3-yl)-allyl	F	F	H	H
I-749	3-(4-chlorophenyl)-but-2-enyl	F	F	H	H
I-750	3-(4-chlorophenyl)-3-fluoro-allyl	F	F	H	H
I-751	3-chloro-4-fluoro-cinnamyl	F	F	H	H

I-752	3,5-dichloro-cinnamyl	F	F	H	H
I-753	5-phenyl-penta-2,4-dienyl	F	F	H	H
I-754	4-isopropylloxycarbonylamino-cinnamyl	F	F	H	H
I-755	3-naphthalen-2-yl-allyl	F	F	H	H
I-756	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	F	F	H	H
I-757	3-(5-chloro-pyridin-2-yl)-allyl	F	F	H	H
I-758	3-pyridin-4-yl-allyl	F	F	H	H
I-759	3-(2-Chloro-pyridin-4-yl)-allyl	F	F	H	H
I-760	4-chlorobenzyl	Cl	F	H	H
I-761	Cinnamyl	Cl	F	H	H
I-762	4-chlorocinnamyl	Cl	F	H	H
I-763	4-fluorocinnamyl	Cl	F	H	H
I-764	4-bromocinnamyl	Cl	F	H	H
I-765	4-trifluoromethylcinnamyl	Cl	F	H	H
I-766	4-trifluoromethoxycinnamyl	Cl	F	H	H
I-767	4-pentafluoroethoxycinnamyl	Cl	F	H	H
I-768	4-methoxycinnamyl	Cl	F	H	H
I-769	4-ethoxycinnamyl	Cl	F	H	H
I-770	4-cyanocinnamyl	Cl	F	H	H
I-771	3-(6-chloro-pyridin-3-yl)-allyl	Cl	F	H	H
I-772	3-(4-chlorophenyl)-but-2-enyl	Cl	F	H	H
I-773	3-(4-chlorophenyl)-3-fluoro-allyl	Cl	F	H	H
I-774	3-chloro-4-fluoro-cinnamyl	Cl	F	H	H
I-775	3,5-dichloro-cinnamyl	Cl	F	H	H
I-776	5-phenyl-penta-2,4-dienyl	Cl	F	H	H
I-777	4-isopropylloxycarbonylamino-cinnamyl	Cl	F	H	H
I-778	3-naphthalen-2-yl-allyl	Cl	F	H	H
I-779	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	Cl	F	H	H
I-780	3-(5-chloro-pyridin-2-yl)-allyl	Cl	F	H	H


I-781	3-pyridin-4-yl-allyl	Cl	F	H	H
I-782	3-(2-Chloro-pyridin-4-yl)-allyl	Cl	F	H	H

Table II provides 782 compounds of formula Ib




wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table III provides 782 compounds of formula Ic



wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1


Table IV provides 782 compounds of formula Id



(Id)

wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1


Table V provides 782 compounds of formula Ie



(Ie)

wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1


Table VI provides 782 compounds of formula If



(If)


wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1

Table VII provides 782 compounds of formula Ig




wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table VIII provides 782 compounds of formula Ih




wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1

Table IX provides 782 compounds of formula Ii




wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table X provides 782 compounds of formula Ij




5 wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XI provides 782 compounds of formula Ik




10 wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XII provides 782 compounds of formula II




wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

15 Table XIII provides 782 compounds of formula Im




wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XIV provides 782 compounds of formula In




wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XV provides 782 compounds of formula Io




wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XVI provides 782 compounds of formula Ip




wherein p is 1, q is 2 and the values of R⁸, R^{4a}, R^{4b}, R^{4c} and R^{4d} are given in Table 1.

Table XVII provides 782 compounds of formula Iq




wherein p is 1, q is 2 and the values of R⁸, R^{4a}, R^{4b}, R^{4c} and R^{4d} are given in Table 1.

Table XVIII provides 782 compounds of formula Ir




wherein p is 1, q is 2 and the values of R⁸, R^{4a}, R^{4b}, R^{4c} and R^{4d} are given in Table 1.

Table XIX provides 782 compounds of formula Is




wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XX provides 782 compounds of formula It



wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXI provides 782 compounds of formula Iu



wherein p is 1, q is 2 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXII provides 782 compounds of formula Ia wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXIII provides 782 compounds of formula Ib wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

5 Table XXIV provides 782 compounds of formula Ic wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXV provides 782 compounds of formula Id wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

10 Table XXVI provides 782 compounds of formula Ie wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXVII provides 782 compounds of formula If wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXVIII provides 782 compounds of formula Ig wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

15 Table XXIX provides 782 compounds of formula Ih wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXX provides 782 compounds of formula Ii wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

20 Table XXXI provides 782 compounds of formula Ij wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXXII provides 782 compounds of formula Ik wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXXIII provides 782 compounds of formula Il wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

25 Table XXXIV provides 782 compounds of formula Im wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXXV provides 782 compounds of formula In wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

30 Table XXXVI provides 782 compounds of formula Io wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXXVII provides 782 compounds of formula Ip wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XXXVIII provides 782 compounds of formula Iq wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

5 Table XXXIX provides 782 compounds of formula Ir wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XLX provides 782 compounds of formula Is wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

10 Table XLI provides 782 compounds of formula It wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XLII provides 782 compounds of formula Iu wherein p is 1, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XLIII provides 782 compounds of formula Ia wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

15 Table XLIV provides 782 compounds of formula Ib wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XLV provides 782 compounds of formula Ic wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

20 Table XLV provides 782 compounds of formula Id wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XLVII provides 782 compounds of formula Ie wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table XLVIII provides 782 compounds of formula If wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

25 Table IL provides 782 compounds of formula Ig wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table L provides 782 compounds of formula Ih wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

30 Table LI provides 782 compounds of formula Ii wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table LII provides 782 compounds of formula Ij wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table LIII provides 782 compounds of formula Ik wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

5 Table LIV provides 782 compounds of formula Il wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table LV provides 782 compounds of formula Im wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

10 Table LVI provides 782 compounds of formula In wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table LVII provides 782 compounds of formula Io wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table LVIII provides 782 compounds of formula Ip wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

15 Table LIX provides 782 compounds of formula Iq wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table LX provides 782 compounds of formula Ir wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

20 Table LXI provides 782 compounds of formula Is wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table LXII provides 782 compounds of formula It wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

Table LXIII provides 782 compounds of formula Iu wherein p is 2, q is 3 and the values of R^8 , R^{4a} , R^{4b} , R^{4c} and R^{4d} are given in Table 1.

25 Mass spectra data were obtained for selected compounds of Tables I to LXIII using LCMS: LC5: 254nm - gradient 10% A to 100% B A=H₂O+0.01%HCOOH
B=CH₃CN/CH₃OH+0.01%HCOOH positive electrospray 150-1000 m/z.


The data are shown in Table 2.

Table 2

Compound	LCMS (Ret. Time, min)	MS data
III-3		464/466
III-49	2'46	498/500/502
XXIV-3	2'38	478/480
XXIV-26	2'49	496/498
XLV.49	2'39	526/528/530
XLVII.49	2'55	560/562/564

The compounds of the invention may be made by a variety of methods. For example
5 they may be prepared according to the reactions of Scheme 1.

SCHEME 1



Thus a compound of formula 1 may be synthesised from compounds of formula 4 by
 5 reaction with an alkylating agent of the formula R8-L, where L is chloride, bromide, iodide
 or a sulfonate (e.g. mesylate or tosylate) or similar leaving group at a temperature of between
 ambient temperature and 100°C, typically ambient temperature, in an organic solvent such as
 acetonitrile, dimethylformamide, dichloromethane, chloroform or 1,2-dichloroethane in the
 presence of a tertiary amine base such as triethylamine or diisopropylethylamine and

optionally catalysed by halide salts such as sodium iodide, potassium iodide or tetrabutylammonium iodide.

Alternatively, a compound of formula 4 may be reacted with an aldehyde of the formula $RCHO$ at a temperature between ambient temperature and $100^{\circ}C$ in an organic solvent such as tetrahydrofuran or ethanol or mixtures of solvents in the presence of a reducing agent such as borane-pyridine complex, sodium borohydride, sodium (triacetoxy)borohydride, sodium cyanoborohydride or such like, to produce a compound of formula 1 where R_8 is CH_2-R .

Compounds of formula 1 may also be obtained from compounds of formula 2b by reaction with a suitable electrophilic species. Compounds of formula 1 where Y is a carbonyl group may be formed by the reaction of compounds of formula 4 with a carboxylic acid derivative of formula $R_1-C(O)-Z$ where Z is chloride, hydroxy, alkoxy or acyloxy at a temperature between $0^{\circ}C$ and $150^{\circ}C$ optionally in an organic solvent such as dichloromethane, chloroform or 1,2-dichloroethane, optionally in the presence of a tertiary amine base such as triethylamine or diisopropylethylamine and optionally in the presence of a coupling agent such as dicyclohexylcarbodiimide. Compounds of formula 1 where Y is a carbonyl group and R_1 is an amino substituent of formula $R'-NH-$ may be formed by the reaction of compounds of formula 2b with an isocyanate of formula $R'-N=C=O$ under similar conditions. Compounds of formula 1 where Y is a group of formula $S(O)_q$ may be formed from compounds of formula 2b by treatment with compounds of formula $R_1-S(O)_q-Cl$ under similar conditions. Compounds of formula 1 where Y is a thiocarbonyl group and R_1 is an amino substituent of formula $R'-NH-$ may be formed by the reaction of compounds of formula 2b with an isothiocyanate of formula $R'-N=C=S$ under similar conditions.

Alternatively compounds of formula 1 where Y is a thiocarbonyl group and R_1 is a carbon substituent may be formed by treatment of compounds of formula 1 where Y is a carbonyl group and R_1 is a carbon substituent with a suitable thionating agent such as Lawesson's reagent.

In the above procedures, acid derivatives of the formula $R_1-C(O)-Z$, isocyanates of formula $R'-N=C=O$, isothiocyanates of formula $R'-N=C=S$ and sulfur electrophiles of formula $R_1-S(O)_q-Cl$ are either known compounds or may be formed from known compounds by known methods by a person skilled in the art.

A compound of formula 4 may be obtained from a compound of formula 3 by reaction with an acid such as trifluoroacetic acid at ambient temperature in an organic solvent such as dichloromethane, chloroform or 1,2-dichloroethane followed by neutralisation of the reaction mixture with an aqueous solution of an inorganic base such as sodium carbonate, sodium bicarbonate or similar compound.

A compound of formula 3 may be obtained from a compound of formula 2a by reaction with a suitable electrophilic species, as described above.

Compounds of formula 2a and 2b may be synthesised as described in Scheme 2. Thus, compounds of formula 2a may be obtained by reacting compounds of formula 6a with compounds of formula 5 at a temperature of between 0°C and 100°C in an organic solvent such as dichloromethane, chloroform or 1,2-dichloroethane in the presence of an acid such as hydrochloric acid or trifluoroacetic acid and optionally a co-solvent such as water, methanol or ethanol. The intermediates formed are subsequently treated with a reducing agent such as sodium borohydride, sodium (triacetoxy)borohydride, sodium cyanoborohydride, triethylsilane or similar at ambient temperature in organic solvent such as ethanol or chloroform.


Similarly, compounds of formula 2b may be synthesised by reacting compounds of formula 6b with compounds of formula 5 using the conditions described above.

Compounds of formula 6a may be obtained from compounds of formula 7a by reaction with a 1-alkoxy substituted phosphonium salt such as methoxymethyl(triphenyl)phosphonium chloride and a base such as potassium *tert*-butoxide at a temperature of 0°C to room temperature in tetrahydrofuran.

Similarly, compounds of formula 6b may be synthesised from compounds of formula 7b using the conditions described above.

Compounds of formula 7a and 7b are either known compounds or may be formed from known compounds by known methods by a person skilled in the art.

SCHEME 2



A compound of formula 2a, in which $p = 1$ and $q = 3$ may be obtained as shown in Scheme 3.

5

Thus, a compound of formula 8 may be reduced to a compound of formula 2b in the presence of a reducing agent such as lithium aluminium, bis(2-methoxyethoxy)aluminium hydride or borane at a temperature of between 0°C and 120°C in an organic solvent such as

tetrahydrofuran, diethyl ether, benzene or toluene. A basic procedure is described in *Synth. Commun.* 1992, 22(5), 729-733.


5 Compounds of formula 8 may be synthesised by cyclising compounds of formula 9 under radical conditions such as tributyltin hydride in the presence of a radical initiator such as 1,1'-azobis(cyclohexanecarbonitrile) in an organic solvent such as benzene or toluene at a temperature of between 60°C and 120°C, followed by removal of the acetyl protecting group using a base such as sodium hydroxide or potassium hydroxide in an organic solvent such as methanol, ethanol or water at a temperature of between 0°C to 100°C.

10 Compounds of formula 9 may be synthesised by acylating compounds of formula 11 using known methods by the person skilled in the art.

Alternatively a compound of formula 8 may be obtained by hydrogenation of a compound of formula 10, which may be obtained from a compound of formula 11 by cyclising a compound of formula 10 under Heck conditions in the presence of a catalyst such as palladium(II) acetate, optionally a ligand such as triphenylphosphine or/and an additive
15 such as tetrabutylammonium bromide and a base such as triethylamine in an organic solvent such as tetrahydrofuran, acetonitrile or dimethylformamide at a temperature of between 50°C to 140°C. A basic procedure is described in WO 01/05790.

Compounds of formula 11 may be synthesised by reacting the known compound of formula 12 (*Chem. Commun.* 1999, 1757-1758) with compounds of formula 13 at
20 temperatures of between 0°C to 60°C in an organic solvent such as dichloromethane, benzene or toluene in the presence of a trialkylaluminium reagent such as trimethylaluminium.

Compounds of formula 13 are either known compounds or may be formed from known compounds by known methods by a person skilled in the art.



Alternatively a compound of formula 2b, in which p = 1 and q = 2 may be obtained as shown in Scheme 4.

Thus, compounds of formula 14 may react with a reducing agent such as bis(2-methoxyethoxy)aluminium hydride at a temperature of between 0°C and 120°C in an organic solvent such as benzene or toluene to provide compounds of formula 2b, in which p = 1 and q = 2.

Compounds of formula 14 may be synthesised from compounds of formula 15 by reaction with an alkylating agent of the formula R8-L, where L is chloride, bromide, iodide or a sulfonate (e.g. mesylate or tosylate) or similar leaving group, as described above.


5 Compounds of formula 15 may be obtained by radical cyclisation of compounds of formula 16 using the method described in *Org. Lett.* 2000, 23, 3599-3601.

Compounds of formula 16 may be synthesised by coupling compounds of formula 18 with the known alcohol 19 (*J. Org. Chem.* 2001, 66, 5545-5551) under Mitsunobu conditions.

10 Compounds of formula 18 are either known compounds or may be formed from known compounds such as 20 by known methods by a person skilled in the art.

Certain compounds of formula 2b and 4 are novel and as such form a further aspect of the invention.

SCHEME 4



The compounds of formula (I) can be used to combat and control infestations of insect pests such as Lepidoptera, Diptera, Hemiptera, Thysanoptera, Orthoptera, Dictyoptera, Coleoptera, Siphonaptera, Hymenoptera and Isoptera and also other invertebrate pests, for example, acarine, nematode and mollusc pests. Insects, acarines, nematodes and molluscs are hereinafter collectively referred to as pests. The pests which may be combated and

controlled by the use of the invention compounds include those pests associated with agriculture (which term includes the growing of crops for food and fibre products), horticulture and animal husbandry, companion animals, forestry and the storage of products of vegetable origin (such as fruit, grain and timber); those pests associated with the damage
 5 of man-made structures and the transmission of diseases of man and animals; and also nuisance pests (such as flies).

Examples of pest species which may be controlled by the compounds of formula (I) include: *Myzus persicae* (aphid), *Aphis gossypii* (aphid), *Aphis fabae* (aphid), *Lygus* spp. (capsids), *Dysdercus* spp. (capsids), *Nilaparvata lugens* (planthopper), *Nephotettix inciticeps*
 10 (leafhopper), *Nezara* spp. (stinkbugs), *Euschistus* spp. (stinkbugs), *Leptocorisa* spp. (stinkbugs), *Frankliniella occidentalis* (thrip), *Thrips* spp. (thrips), *Leptinotarsa decemlineata* (Colorado potato beetle), *Anthonomus grandis* (boll weevil), *Aonidiella* spp. (scale insects), *Trialeurodes* spp. (white flies), *Bemisia tabaci* (white fly), *Ostrinia nubilalis* (European corn borer), *Spodoptera littoralis* (cotton leafworm), *Heliothis virescens* (tobacco
 15 budworm), *Helicoverpa armigera* (cotton bollworm), *Helicoverpa zea* (cotton bollworm), *Sylepta derogata* (cotton leaf roller), *Pieris brassicae* (white butterfly), *Plutella xylostella* (diamond back moth), *Agrotis* spp. (cutworms), *Chilo suppressalis* (rice stem borer), *Locusta migratoria* (locust), *Chortiocetes terminifera* (locust), *Diabrotica* spp. (rootworms), *Panonychus ulmi* (European red mite), *Panonychus citri* (citrus red mite), *Tetranychus*
 20 *urticae* (two-spotted spider mite), *Tetranychus cinnabarinus* (carmine spider mite), *Phyllocoptruta oleivora* (citrus rust mite), *Polyphagotarsonemus latus* (broad mite), *Brevipalpus* spp. (flat mites), *Boophilus microplus* (cattle tick), *Dermacentor variabilis* (American dog tick), *Ctenocephalides felis* (cat flea), *Liriomyza* spp. (leafminer), *Musca domestica* (housefly), *Aedes aegypti* (mosquito), *Anopheles* spp. (mosquitoes), *Culex* spp.
 25 (mosquitoes), *Lucillia* spp. (blowflies), *Blattella germanica* (cockroach), *Periplaneta americana* (cockroach), *Blatta orientalis* (cockroach), termites of the Mastotermitidae (for example *Mastotermes* spp.), the Kalotermitidae (for example *Neotermes* spp.), the Rhinotermitidae (for example *Coptotermes formosanus*, *Reticulitermes flavipes*, *R. speratu*, *R. virginicus*, *R. hesperus*, and *R. santonensis*) and the Termitidae (for example *Globitermes*
 30 *sulphureus*), *Solenopsis geminata* (fire ant), *Monomorium pharaonis* (pharaoh's ant), *Damalinea* spp. and *Linognathus* spp. (biting and sucking lice), *Meloidogyne* spp. (root knot nematodes), *Globodera* spp. and *Heterodera* spp. (cyst nematodes), *Pratylenchus* spp. (lesion

nematodes), *Rhodopholus* spp. (banana burrowing nematodes), *Tylenchulus* spp. (citrus nematodes), *Haemonchus contortus* (barber pole worm), *Caenorhabditis elegans* (vinegar eelworm), *Trichostrongylus* spp. (gastro intestinal nematodes) and *Deroceras reticulatum* (slug).

5 The invention therefore provides a method of combating and controlling insects, acarines, nematodes or molluscs which comprises applying an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a compound of formula (I), or a composition containing a compound of formula (I), to a pest, a locus of pest, or to a plant susceptible to attack by a pest, The compounds of formula (I) are preferably used against
10 insects, acarines or nematodes.

The term "plant" as used herein includes seedlings, bushes and trees.

In order to apply a compound of formula (I) as an insecticide, acaricide, nematocide or molluscicide to a pest, a locus of pest, or to a plant susceptible to attack by a pest, a compound of formula (I) is usually formulated into a composition which includes, in addition
15 to the compound of formula (I), a suitable inert diluent or carrier and, optionally, a surface active agent (SFA). SFAs are chemicals which are able to modify the properties of an interface (for example, liquid/solid, liquid/air or liquid/liquid interfaces) by lowering the interfacial tension and thereby leading to changes in other properties (for example dispersion, emulsification and wetting). It is preferred that all compositions (both solid and liquid
20 formulations) comprise, by weight, 0.0001 to 95%, more preferably 1 to 85%, for example 5 to 60%, of a compound of formula (I). The composition is generally used for the control of pests such that a compound of formula (I) is applied at a rate of from 0.1g to 10kg per hectare, preferably from 1g to 6kg per hectare, more preferably from 1g to 1kg per hectare.

When used in a seed dressing, a compound of formula (I) is used at a rate of 0.0001g
25 to 10g (for example 0.001g or 0.05g), preferably 0.005g to 10g, more preferably 0.005g to 4g, per kilogram of seed.

In another aspect the present invention provides an insecticidal, acaricidal, nematocidal or molluscicidal composition comprising an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a compound of formula (I) and a
30 suitable carrier or diluent therefor. The composition is preferably an insecticidal, acaricidal, nematocidal or molluscicidal composition.

In a still further aspect the invention provides a method of combating and controlling pests at a locus which comprises treating the pests or the locus of the pests with an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a composition comprising a compound of formula (I). The compounds of formula (I) are preferably used against insects, acarines or nematodes.

The compositions can be chosen from a number of formulation types, including dustable powders (DP), soluble powders (SP), water soluble granules (SG), water dispersible granules (WG), wettable powders (WP), granules (GR) (slow or fast release), soluble concentrates (SL), oil miscible liquids (OL), ultra low volume liquids (UL), emulsifiable concentrates (EC), dispersible concentrates (DC), emulsions (both oil in water (EW) and water in oil (EO)), micro-emulsions (ME), suspension concentrates (SC), aerosols, fogging/smoke formulations, capsule suspensions (CS) and seed treatment formulations. The formulation type chosen in any instance will depend upon the particular purpose envisaged and the physical, chemical and biological properties of the compound of formula (I).

Dustable powders (DP) may be prepared by mixing a compound of formula (I) with one or more solid diluents (for example natural clays, kaolin, pyrophyllite, bentonite, alumina, montmorillonite, kieselguhr, chalk, diatomaceous earths, calcium phosphates, calcium and magnesium carbonates, sulphur, lime, flours, talc and other organic and inorganic solid carriers) and mechanically grinding the mixture to a fine powder.

Soluble powders (SP) may be prepared by mixing a compound of formula (I) with one or more water-soluble inorganic salts (such as sodium bicarbonate, sodium carbonate or magnesium sulphate) or one or more water-soluble organic solids (such as a polysaccharide) and, optionally, one or more wetting agents, one or more dispersing agents or a mixture of said agents to improve water dispersibility/solubility. The mixture is then ground to a fine powder. Similar compositions may also be granulated to form water soluble granules (SG).

Wettable powders (WP) may be prepared by mixing a compound of formula (I) with one or more solid diluents or carriers, one or more wetting agents and, preferably, one or more dispersing agents and, optionally, one or more suspending agents to facilitate the dispersion in liquids. The mixture is then ground to a fine powder. Similar compositions may also be granulated to form water dispersible granules (WG).

Granules (GR) may be formed either by granulating a mixture of a compound of formula (I) and one or more powdered solid diluents or carriers, or from pre-formed blank

granules by absorbing a compound of formula (I) (or a solution thereof, in a suitable agent) in a porous granular material (such as pumice, attapulgite clays, fuller's earth, kieselguhr, diatomaceous earths or ground corn cobs) or by adsorbing a compound of formula (I) (or a solution thereof, in a suitable agent) on to a hard core material (such as sands, silicates, mineral carbonates, sulphates or phosphates) and drying if necessary. Agents which are commonly used to aid absorption or adsorption include solvents (such as aliphatic and aromatic petroleum solvents, alcohols, ethers, ketones and esters) and sticking agents (such as polyvinyl acetates, polyvinyl alcohols, dextrans, sugars and vegetable oils). One or more other additives may also be included in granules (for example an emulsifying agent, wetting agent or dispersing agent).

Dispersible Concentrates (DC) may be prepared by dissolving a compound of formula (I) in water or an organic solvent, such as a ketone, alcohol or glycol ether. These solutions may contain a surface active agent (for example to improve water dilution or prevent crystallisation in a spray tank).

Emulsifiable concentrates (EC) or oil-in-water emulsions (EW) may be prepared by dissolving a compound of formula (I) in an organic solvent (optionally containing one or more wetting agents, one or more emulsifying agents or a mixture of said agents). Suitable organic solvents for use in ECs include aromatic hydrocarbons (such as alkylbenzenes or alkylnaphthalenes, exemplified by SOLVESSO 100, SOLVESSO 150 and SOLVESSO 200; SOLVESSO is a Registered Trade Mark), ketones (such as cyclohexanone or methylcyclohexanone) and alcohols (such as benzyl alcohol, furfuryl alcohol or butanol), N-alkylpyrrolidones (such as N-methylpyrrolidone or N-octylpyrrolidone), dimethyl amides of fatty acids (such as C₈-C₁₀ fatty acid dimethylamide) and chlorinated hydrocarbons. An EC product may spontaneously emulsify on addition to water, to produce an emulsion with sufficient stability to allow spray application through appropriate equipment. Preparation of an EW involves obtaining a compound of formula (I) either as a liquid (if it is not a liquid at room temperature, it may be melted at a reasonable temperature, typically below 70°C) or in solution (by dissolving it in an appropriate solvent) and then emulsifying the resultant liquid or solution into water containing one or more SFAs, under high shear, to produce an emulsion. Suitable solvents for use in EWs include vegetable oils, chlorinated hydrocarbons (such as chlorobenzenes), aromatic solvents (such as alkylbenzenes or alkylnaphthalenes) and other appropriate organic solvents which have a low solubility in water.

Microemulsions (ME) may be prepared by mixing water with a blend of one or more solvents with one or more SFAs, to produce spontaneously a thermodynamically stable isotropic liquid formulation. A compound of formula (I) is present initially in either the water or the solvent/SFA blend. Suitable solvents for use in MEs include those hereinbefore described for use in ECs or in EWs. An ME may be either an oil-in-water or a water-in-oil system (which system is present may be determined by conductivity measurements) and may be suitable for mixing water-soluble and oil-soluble pesticides in the same formulation. An ME is suitable for dilution into water, either remaining as a microemulsion or forming a conventional oil-in-water emulsion.

Suspension concentrates (SC) may comprise aqueous or non-aqueous suspensions of finely divided insoluble solid particles of a compound of formula (I). SCs may be prepared by ball or bead milling the solid compound of formula (I) in a suitable medium, optionally with one or more dispersing agents, to produce a fine particle suspension of the compound. One or more wetting agents may be included in the composition and a suspending agent may be included to reduce the rate at which the particles settle. Alternatively, a compound of formula (I) may be dry milled and added to water, containing agents hereinbefore described, to produce the desired end product.

Aerosol formulations comprise a compound of formula (I) and a suitable propellant (for example *n*-butane). A compound of formula (I) may also be dissolved or dispersed in a suitable medium (for example water or a water miscible liquid, such as *n*-propanol) to provide compositions for use in non-pressurised, hand-actuated spray pumps.

A compound of formula (I) may be mixed in the dry state with a pyrotechnic mixture to form a composition suitable for generating, in an enclosed space, a smoke containing the compound.

Capsule suspensions (CS) may be prepared in a manner similar to the preparation of EW formulations but with an additional polymerisation stage such that an aqueous dispersion of oil droplets is obtained, in which each oil droplet is encapsulated by a polymeric shell and contains a compound of formula (I) and, optionally, a carrier or diluent therefor. The polymeric shell may be produced by either an interfacial polycondensation reaction or by a coacervation procedure. The compositions may provide for controlled release of the compound of formula (I) and they may be used for seed treatment. A compound of formula

(I) may also be formulated in a biodegradable polymeric matrix to provide a slow, controlled release of the compound.

A composition may include one or more additives to improve the biological performance of the composition (for example by improving wetting, retention or distribution on surfaces; resistance to rain on treated surfaces; or uptake or mobility of a compound of formula (I)). Such additives include surface active agents, spray additives based on oils, for example certain mineral oils or natural plant oils (such as soy bean and rape seed oil), and blends of these with other bio-enhancing adjuvants (ingredients which may aid or modify the action of a compound of formula (I)).

A compound of formula (I) may also be formulated for use as a seed treatment, for example as a powder composition, including a powder for dry seed treatment (DS), a water soluble powder (SS) or a water dispersible powder for slurry treatment (WS), or as a liquid composition, including a flowable concentrate (FS), a solution (LS) or a capsule suspension (CS). The preparations of DS, SS, WS, FS and LS compositions are very similar to those of, respectively, DP, SP, WP, SC and DC compositions described above. Compositions for treating seed may include an agent for assisting the adhesion of the composition to the seed (for example a mineral oil or a film-forming barrier).

Wetting agents, dispersing agents and emulsifying agents may be surface SFAs of the cationic, anionic, amphoteric or non-ionic type.

Suitable SFAs of the cationic type include quaternary ammonium compounds (for example cetyltrimethyl ammonium bromide), imidazolines and amine salts.

Suitable anionic SFAs include alkali metals salts of fatty acids, salts of aliphatic monoesters of sulphuric acid (for example sodium lauryl sulphate), salts of sulphonated aromatic compounds (for example sodium dodecylbenzenesulphonate, calcium dodecylbenzenesulphonate, butylnaphthalene sulphonate and mixtures of sodium di-*isopropyl*- and tri-*isopropyl*-naphthalene sulphonates), ether sulphates, alcohol ether sulphates (for example sodium laureth-3-sulphate), ether carboxylates (for example sodium laureth-3-carboxylate), phosphate esters (products from the reaction between one or more fatty alcohols and phosphoric acid (predominately mono-esters) or phosphorus pentoxide (predominately di-esters), for example the reaction between lauryl alcohol and tetraphosphoric acid; additionally these products may be ethoxylated), sulposuccinamates, paraffin or olefine sulphonates, taurates and lignosulphonates.

Suitable SFAs of the amphoteric type include betaines, propionates and glycines.

Suitable SFAs of the non-ionic type include condensation products of alkylene oxides, such as ethylene oxide, propylene oxide, butylene oxide or mixtures thereof, with fatty alcohols (such as oleyl alcohol or cetyl alcohol) or with alkylphenols (such as octylphenol, nonylphenol or octylcresol); partial esters derived from long chain fatty acids or hexitol anhydrides; condensation products of said partial esters with ethylene oxide; block polymers (comprising ethylene oxide and propylene oxide); alkanolamides; simple esters (for example fatty acid polyethylene glycol esters); amine oxides (for example lauryl dimethyl amine oxide); and lecithins.

Suitable suspending agents include hydrophilic colloids (such as polysaccharides, polyvinylpyrrolidone or sodium carboxymethylcellulose) and swelling clays (such as bentonite or attapulgite).

A compound of formula (I) may be applied by any of the known means of applying pesticidal compounds. For example, it may be applied, formulated or unformulated, to the pests or to a locus of the pests (such as a habitat of the pests, or a growing plant liable to infestation by the pests) or to any part of the plant, including the foliage, stems, branches or roots, to the seed before it is planted or to other media in which plants are growing or are to be planted (such as soil surrounding the roots, the soil generally, paddy water or hydroponic culture systems), directly or it may be sprayed on, dusted on, applied by dipping, applied as a cream or paste formulation, applied as a vapour or applied through distribution or incorporation of a composition (such as a granular composition or a composition packed in a water-soluble bag) in soil or an aqueous environment.

A compound of formula (I) may also be injected into plants or sprayed onto vegetation using electrodynamic spraying techniques or other low volume methods, or applied by land or aerial irrigation systems.

Compositions for use as aqueous preparations (aqueous solutions or dispersions) are generally supplied in the form of a concentrate containing a high proportion of the active ingredient, the concentrate being added to water before use. These concentrates, which may include DCs, SCs, ECs, EWs, MEs SGs, SPs, WPs, WGs and CSs, are often required to withstand storage for prolonged periods and, after such storage, to be capable of addition to water to form aqueous preparations which remain homogeneous for a sufficient time to enable them to be applied by conventional spray equipment. Such aqueous preparations may

contain varying amounts of a compound of formula (I) (for example 0.0001 to 10%, by weight) depending upon the purpose for which they are to be used.

A compound of formula (I) may be used in mixtures with fertilisers (for example nitrogen-, potassium- or phosphorus-containing fertilisers). Suitable formulation types include granules of fertiliser. The mixtures suitably contain up to 25% by weight of the compound of formula (I).

The invention therefore also provides a fertiliser composition comprising a fertiliser and a compound of formula (I).

The compositions of this invention may contain other compounds having biological activity, for example micronutrients or compounds having fungicidal activity or which possess plant growth regulating, herbicidal, insecticidal, nematocidal or acaricidal activity.

The compound of formula (I) may be the sole active ingredient of the composition or it may be admixed with one or more additional active ingredients such as a pesticide, fungicide, synergist, herbicide or plant growth regulator where appropriate. An additional active ingredient may: provide a composition having a broader spectrum of activity or increased persistence at a locus; synergise the activity or complement the activity (for example by increasing the speed of effect or overcoming repellency) of the compound of formula (I); or help to overcome or prevent the development of resistance to individual components. The particular additional active ingredient will depend upon the intended utility of the composition. Examples of suitable pesticides include the following:

- a) Pyrethroids, such as permethrin, cypermethrin, fenvalerate, esfenvalerate, deltamethrin, cyhalothrin (in particular lambda-cyhalothrin), bifenthrin, fenpropathrin, cyfluthrin, tefluthrin, fish safe pyrethroids (for example ethofenprox), natural pyrethrin, tetramethrin, s-bioallethrin, fenfluthrin, prallethrin or 5-benzyl-3-furylmethyl-(E)-(1R,3S)-2,2-dimethyl-3-(2-oxothiolan-3-ylidenemethyl)cyclopropane carboxylate;
- b) Organophosphates, such as, profenofos, sulprofos, acephate, methyl parathion, azinphos-methyl, demeton-s-methyl, heptenophos, thiometon, fenamiphos, monocrotophos, profenofos, triazophos, methamidophos, dimethoate, phosphamidon, malathion, chlorpyrifos, phosalone, terbufos, fensulfothion, fonofos, phorate, phoxim, pirimiphos-methyl, pirimiphos-ethyl, fenitrothion, fosthiazate or diazinon;

- c) Carbamates (including aryl carbamates), such as pirimicarb, triazamate, cloethocarb, carbofuran, furathiocarb, ethiofencarb, aldicarb, thiofurox, carbosulfan, bendiocarb, fenobucarb, propoxur, methomyl or oxamyl;
- d) Benzoyl ureas, such as diflubenzuron, triflumuron, hexaflumuron, flufenoxuron or
5 chlorfluazuron;
- e) Organic tin compounds, such as cyhexatin, fenbutatin oxide or azocyclotin;
- f) Pyrazoles, such as tebufenpyrad and fenpyroximate;
- g) Macrolides, such as avermectins or milbemycins, for example abamectin, emamectin benzoate, ivermectin, milbemycin, spinosad or azadirachtin;
- 10 h) Hormones or pheromones;
- i) Organochlorine compounds such as endosulfan, benzene hexachloride, DDT, chlordane or dieldrin;
- j) Amidines, such as chlordimeform or amitraz;
- k) Fumigant agents, such as chloropicrin, dichloropropane, methyl bromide or metam;
- 15 l) Chloronicotinyl compounds such as imidacloprid, thiacloprid, acetamiprid, nitenpyram or thiamethoxam;
- m) Diacylhydrazines, such as tebufenozide, chromafenozide or methoxyfenozide;
- n) Diphenyl ethers, such as diofenolan or pyriproxifen;
- o) Indoxacarb;
- 20 p) Chlorfenapyr; or
- q) Pymetrozine.

In addition to the major chemical classes of pesticide listed above, other pesticides having particular targets may be employed in the composition, if appropriate for the intended utility of the composition. For instance, selective insecticides for particular crops, for
25 example stemborer specific insecticides (such as cartap) or hopper specific insecticides (such as buprofezin) for use in rice may be employed. Alternatively insecticides or acaricides specific for particular insect species/stages may also be included in the compositions (for example acaricidal ovo-larvicides, such as clofentezine, flubenzimine, hexythiazox or tetradifon; acaricidal motilicides, such as dicofol or propargite; acaricides, such as
30 bromopropylate or chlorobenzilate; or growth regulators, such as hydramethylnon, cyromazine, methoprene, chlorfluazuron or diflubenzuron).

Examples of fungicidal compounds which may be included in the composition of the invention are (*E*)-*N*-methyl-2-[2-(2,5-dimethylphenoxy)methyl]phenyl]-2-methoxy-iminoacetamide (SSF-129), 4-bromo-2-cyano-*N,N*-dimethyl-6-trifluoromethylbenzimidazole-1-sulphonamide, α -[*N*-(3-chloro-2,6-xylyl)-2-methoxyacetamido]- γ -butyrolactone, 4-chloro-2-cyano-*N,N*-dimethyl-5-*p*-tolylimidazole-1-sulfonamide (IKF-916, cyamidazosulfamid), 3-5-dichloro-*N*-(3-chloro-1-ethyl-1-methyl-2-oxopropyl)-4-methylbenzamide (RH-7281, zoxamide), *N*-allyl-4,5,-dimethyl-2-trimethylsilylthiophene-3-carboxamide (MON65500), *N*-(1-cyano-1,2-dimethylpropyl)-2-(2,4-dichlorophenoxy)propionamide (AC382042), *N*-(2-methoxy-5-pyridyl)-cyclopropane carboxamide, acibenzolar (CGA245704), alanycarb, aldimorph, anilazine, azaconazole, azoxystrobin, benalaxyl, benomyl, biloxazol, bitertanol, blasticidin S, bromuconazole, bupirimate, captafol, captan, carbendazim, carbendazim chlorhydrate, carboxin, carpropamid, carvone, CGA41396, CGA41397, chinomethionate, chlorothalonil, chlorozoline, clozylacon, copper containing compounds such as copper oxychloride, copper oxyquinolate, copper sulphate, copper tallate and Bordeaux mixture, cymoxanil, cyproconazole, cyprodinil, debacarb, di-2-pyridyl disulphide 1,1'-dioxide, dichlofluanid, diclomezine, dicloran, diethofencarb, difenoconazole, difenzoquat, diflumetorim, *O,O*-di-*iso*-propyl-*S*-benzyl thiophosphate, dimefluazole, dimetconazole, dimethomorph, dimethirimol, diniconazole, dinocap, dithianon, dodecyl dimethyl ammonium chloride, dodemorph, dodine, doguadine, edifenphos, epoxiconazole, ethirimol, ethyl(*Z*)-*N*-benzyl-*N*-([methyl(methyl-thioethylideneaminooxycarbonyl)amino]thio)- β -alaninate, etridiazole, famoxadone, fenamidone (RPA407213), fenarimol, fenbuconazole, fenfuram, fenhexamid (KBR2738), fencpiclonil, fenpropidin, fenpropimorph, fentin acetate, fentin hydroxide, ferbam, ferimzone, fluazinam, fludioxonil, flumetover, fluoroimide, fluquinconazole, flusilazole, flutolanil, flutriafol, folpet, fuberidazole, furalaxyl, furametpyr, guazatine, hexaconazole, hydroxyisoxazole, hymexazole, imazalil, imibenconazole, iminoctadine, iminoctadine triacetate, ipconazole, iprobenfos, iprodione, iprovalicarb (SZX0722), isopropanyl butyl carbamate, isoprothiolane, kasugamycin, kresoxim-methyl, LY186054, LY211795, LY248908, mancozeb, maneb, mefenoxam, mepanipyrim, mepronil, metalaxyl, metconazole, metiram, metiram-zinc, metominostrobin, myclobutanil, neoasozin, nickel dimethyldithiocarbamate, nitrothal-*isopropyl*, nuarimol, ofurace, organomercury compounds, oxadixyl, oxasulfuron, oxolinic acid, oxpoconazole, oxycarboxin, pefurazoate, penconazole, pencycuron, phenazin oxide, phosetyl-A1, phosphorus acids, phthalide,

picoxystrobin (ZA1963), polyoxin D, polyram, probenazole, prochloraz, procymidone, propamocarb, propiconazole, propineb, propionic acid, pyrazophos, pyrifenoxy, pyrimethanil, pyroquilon, pyroxyfur, pyrrolnitrin, quaternary ammonium compounds, quinomethionate, quinoxifen, quintozone, sipconazole (F-155), sodium pentachlorophenate, spiroxamine, streptomycin, sulphur, tebuconazole, tecloftalam, tecnazene, tetraconazole, thiabendazole, thifluzamid, 2-(thiocyanomethylthio)benzothiazole, thiophanate-methyl, thiram, timibenconazole, tolclofos-methyl, tolylfluanid, triadimefon, triadimenol, triazbutyl, triazoxide, tricyclazole, tridemorph, trifloxystrobin (CGA279202), triforine, triflumizole, triticonazole, validamycin A, vapam, vinclozolin, zineb and ziram.

10 The compounds of formula (I) may be mixed with soil, peat or other rooting media for the protection of plants against seed-borne, soil-borne or foliar fungal diseases.

Examples of suitable synergists for use in the compositions include piperonyl butoxide, sesamex, safroxan and dodecyl imidazole.

15 Suitable herbicides and plant-growth regulators for inclusion in the compositions will depend upon the intended target and the effect required.


An example of a rice selective herbicide which may be included is propanil. An example of a plant growth regulator for use in cotton is PIX™.

20 Some mixtures may comprise active ingredients which have significantly different physical, chemical or biological properties such that they do not easily lend themselves to the same conventional formulation type. In these circumstances other formulation types may be prepared. For example, where one active ingredient is a water insoluble solid and the other a water insoluble liquid, it may nevertheless be possible to disperse each active ingredient in the same continuous aqueous phase by dispersing the solid active ingredient as a suspension (using a preparation analogous to that of an SC) but dispersing the liquid active ingredient as an emulsion (using a preparation analogous to that of an EW). The resultant composition is a
25 suspoemulsion (SE) formulation.

The invention is illustrated by the following Examples:

EXAMPLE 1

This Example illustrates the preparation of compound XLV.49, 5-chloro-1-(2-chloropyridin-4-yl)carbonyl-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[perhydro-azepine-3,3'-piperidine]



Step A: 4-Perhydroazepinone hydrochloride (2.5 g, prepared according to Synth. Commun. 1992, 1249-1258) was suspended in acetonitrile (80 ml); diisopropylethylamine (4.4 ml) and 4-chlorocinnamyl chloride (2.7 g) were successively added at room temperature and the resulting reaction mixture was stirred at room temperature for 20 hours. The solvent was removed *in vacuo* and the residue purified by silica gel chromatography (eluent cyclohexane:ethyl acetate 6:4) to afford 1-(*trans*-3-(4-chlorophenyl)allyl)-perhydroazepin-4-one (1.84 g) as a foam. MS (ES+) 264/266 (M+H⁺).

Step B: To a stirred suspension of methoxymethyltriphenylphosphonium chloride (3.9 g) in tetrahydrofuran (30 ml) at 0°C under argon was added portionwise potassium tert-butoxide (1.3 g) over 30 min. 1-(*trans*-3-(4-chlorophenyl)allyl)-perhydroazepin-4-one (1.5 g) dissolved in a minimum volume of tetrahydrofuran was added to the resulting orange solution and the resulting mixture was stirred at room temperature for 2 hours, quenched by addition of water, extracted twice with ethyl acetate, the organic layers were dried (sodium sulphate) and concentrated *in vacuo*. Silica gel chromatography (eluent cyclohexane:ethyl acetate 7:3) of the residue afforded 1-[(E)-3-(4-Chloro-phenyl)-allyl]-4-[1-methoxy-meth-

(Z)-ylidene]-perhydro-azepine (1.1 g) as an oil (1:1 mixture of isomers). MS (ES+) 292/296 (M+H⁺).


Step C: A mixture of -[(E)-3-(4-Chloro-phenyl)-allyl]-4-[1-methoxy-meth-(Z)-ylidene]-perhydro-azepine (0.6 g) and 4-chlorophenylhydrazine hydrochloride (0.41 g) in
5 chloroform (20 ml) was treated with trifluoroacetic acid (2.1 ml) and heated at reflux under argon for 18 hours. The reaction mixture was cooled to room temperature, triethylsilane (3.1 ml) was added and the solution refluxed for 2 hours. The reaction mixture was cooled to room temperature, diluted with dichloromethane, neutralised with 30% aqueous ammonium hydroxide, washed with brine, dried (sodium sulphate) and concentrated. The dark residue
10 was purified by silica gel chromatography (cyclohexane:ethyl acetate 1:9) to afford 5-chloro-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[perhydro-azepine-3,3'-piperidine] (529 mg). MS (ES+) 387/389 (M+H⁺).

Step D: To a solution of 5-chloro-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[perhydro-azepine-3,3'-piperidine] obtained in Step C (250 mg) and triethylamine (0.42 ml) in
15 dichloromethane (10 ml) at 0°C was added 2-chloro-isonicotinoyl chloride (220 mg) and the resulting solution was kept at room temperature for 18 hours., diluted with dichloromethane, washed with diluted aqueous sodium bicarbonate, dried (sodium sulphate) and concentrated. Silica gel chromatography of the residue (cyclohexane:ethyl acetate 3:7) afforded the title compound as a yellow solid (181 mg); M.p. 86-90°C; MS (ES+) 526/528/530 (M+H⁺)

20 Compound XLV11.49 (M.p. 166-170°C) was prepared according to procedures analogous to those described in Example 1.

EXAMPLE 2

25 This Example illustrates the preparation of compound XXIV.3, 1-(2-chloropyridin-4-yl)carbonyl-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indoline-3,3'-piperidine]



Step A: Trimethylaluminium (2M in heptane, 3 ml) was added dropwise to a solution of 2-bromoaniline (860 mg) in dichloromethane (15 ml). After gas evolution ceased, 5,6-dihydro-2H-pyridine-1,3-dicarboxylic acid 1-tert-butyl ester 3-methyl ester (*Chem. Commun.* 1999, 1757-1758, 1.2 g) dissolved in dichloromethane (10 ml) was added and the resulting mixture was refluxed for 5 hrs. The solution was cooled to 0°C, quenched by careful addition of saturated aqueous sodium bicarbonate (10 ml) and extracted with dichloromethane. The organic layer was washed with brine, dried (Na₂SO₄) and concentrated in vacuo. The residue was purified by silica gel chromatography (eluent cyclohexane:ethyl acetate 85:15) to afford 5-(2-bromo-phenylcarbamoyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester (1.2 g), which was characterised by its mass and NMR spectra.

Step B: To a stirred solution of the compound obtained in Step A (3.24 g) in dimethylformamide (60 ml) under argon were added successively triethylamine (3 ml), tetrabutylammonium bromide (3.2 g) and palladium(II) acetate (386 mg) and the resulting mixture was heated under reflux for 5 hours, cooled to room temperature, poured into brine and extracted with ethyl acetate. The organic layer was washed with HCl 1N then water, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel chromatography (eluent cyclohexane:ethyl acetate 8:2) to afford spiro[indolin-2-one-3,3'-(1',2',3',4'-tetrahydropyridine)]-1'-carboxylic acid tert-butyl ester (1.2 g), which was characterised by its mass and NMR spectra.

Step C: Spiro[indolin-2-one-3,3'-piperidine]-1'-carboxylic acid tert-butyl ester obtained in Step B (1.12 g) was hydrogenated (1 atm.) in 15 ml tetrahydrofuran in the presence of 10% Pd/C (0.6 g) to afford after standard work-up spiro[indolin-2-one-3,3'-piperidine]-1'-carboxylic acid tert-butyl ester (1.06 g), which was characterised by its mass and NMR spectra. MS (ES⁺) 203 (M-CO₂-isoprene+H⁺), 247 (M-isoprene+H⁺), 303 (M+H⁺).


Step D: A solution of spiro[indolin-2-one-3,3'-piperidine]-1'-carboxylic acid tert-butyl ester obtained in Step C (650 mg) in toluene (20 ml) at 70°C under argon was treated with sodium bis(2-methoxyethoxy)aluminium hydride (Red-Al, 65% in toluene, 1.3 ml) and the reaction mixture was stirred at 75°C for 2 hours, cooled to room temperature, quenched by addition of ethyl acetate (20 ml), stirred for 15 min. and concentrated *in vacuo*. The residue was diluted with ethyl acetate, washed with water and brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel chromatography (eluent cyclohexane:ethyl acetate 7:3) to afford spiro[indoline-3,3'-piperidine]-1'-carboxylic acid tert-butyl ester (331 mg) as a colorless oil, which was characterised by its mass and NMR spectra. MS (ES+) 233 (M-isoprene + H⁺), 289 (M+H⁺).

Step E: To a solution of spiro[indoline-3,3'-piperidine]-1'-carboxylic acid tert-butyl ester obtained in Step D (140 mg) and triethylamine (0.28 ml) in dichloromethane (5 ml) at 0°C was added 2-chloroisonicotinoyl chloride (176 mg). After stirring at room temperature for 1 hour, the solution was diluted with dichloromethane, washed with water, dried (sodium sulfate and concentrated *in vacuo*. The residue was dissolved in dichloromethane (12.5 ml) and trifluoroacetic acid (1.25 ml) was added. The reaction mixture was stirred at room temperature for 1 hour, diluted with dichloromethane, neutralised with saturated sodium bicarbonate, dried (sodium sulfate) and concentrated *in vacuo*. The residue was dissolved in acetonitrile (10 ml); diisopropylethylamine (0.13 ml) and 4-chlorocinnamyl chloride (93 mg) were added and the resulting mixture was stirred at room temperature for 12 hours then the solvent was evaporated *in vacuo*. The residue was purified by silica gel chromatography (eluent cyclohexane:ethyl acetate 6:4) to afford the title product (143 mg), which was characterised by its mass and NMR spectra. MS (ES+) 478/480 (M+H⁺).

Compound XXIV.26 was prepared according to procedures analogous to those described in Example 2.

EXAMPLE 3

This Example illustrates the preparation of compound III.49, 5-Chloro-1-(2-chloropyridin-4-yl)carbonyl-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indoline-3,3'-pyrrolidine]



Step A: To a stirred solution of triphenylphosphine (2.75 g) in tetrahydrofuran (60 ml) at -10°C under argon was added dropwise diisopropylazodicarboxylate (DIAD, 2.1 ml); the resulting suspension was stirred at -10°C for 15 min then 4-chloro-2-iodo-N-methanesulfonyl-aniline (3.1 g) was added as a solid followed by 2-methylene-butane-1,4-diol (0.95 g) dissolved in a minimum of tetrahydrofuran. The reaction mixture was stirred at room temperature for 6 hours, the solvent was evaporated *in vacuo* and the residue was purified by silica gel chromatography (eluent ethyl acetate: cyclohexane 4:6) to afford N-(4-Chloro-2-iodo-phenyl)-N-(4-hydroxy-2-methylene-butyl)-methanesulfonamide (3.5 g) contaminated with triphenylphosphine oxide.

Step B: To a stirred solution of N-(4-Chloro-2-iodo-phenyl)-N-(4-hydroxy-2-methylene-butyl)-methanesulfonamide obtained in Step A and triphenylphosphine (3.5 g) in dimethylacetamide (80 ml) at -10°C under argon was added carbon tetrabromide (4.5 g). The reaction mixture was stirred at -10°C for 45 min (a precipitate formed). Sodium azide (2 g) was added in one portion and the reaction mixture was stirred at 45°C for 1 hour, cooled to room temperature, poured into water, extracted with ethyl acetate; the organic layer was washed with brine, dried (Na_2SO_4) and concentrated *in vacuo*. The residue was purified by silica gel chromatography (eluent cyclohexane:ethyl acetate 7:3) to afford N-(4-Chloro-2-iodo-phenyl)-N-(4-azido-2-methylene-butyl)-methanesulfonamide (1.96 g). ^1H NMR (CDCl_3 , 400 MHz) 2.50 (m, 2H), 3.48 (m, 2H), 4.19 (d, $J = 12.0$ Hz, 1H), 4.40 (d, $J = 12.0$ Hz, 1H), 4.90 (s, 1H), 5.01 (s, 1H), 7.29 (d, $J = 8.5$ Hz, 1H), 7.36 (dd, $J = 0.9$ Hz, 8.5 Hz, 1H), 7.92 (d, $J = 0.9$ Hz, 1H).

Step C: A degassed solution of N-(4-Chloro-2-iodo-phenyl)-N-(4-azido-2-methylene-butyl)-methanesulfonamide (1.38 g) in benzene (200 ml) was heated to reflux under argon. Tris(trimethylsilyl)silane (1.32 ml) was added dropwise followed by 1,1'-azobis(cyclohexane carbonitrile) (110 mg). The reaction mixture was stirred at reflux for 20 hours then

concentrated *in vacuo*. The residue was dissolved in ethyl acetate (60 ml), extracted with HCl 2N (3x60 ml). The aqueous layer was basified with 2N sodium hydroxide (250 ml) then extracted with ethyl acetate (3x150 ml). The combined organic layers were dried over sodium sulfate and the solvent evaporated *in vacuo* to afford 1-methanesulfonyl-5-chloro-
5 spiro[indoline-3,3'-pyrrolidine] (766 mg) which was used as such for the next step.

Step D: To a solution of 1-methanesulfonyl-5-chloro-spiro[indoline-3,3'-pyrrolidine] (725 mg) in acetonitrile (40 ml) were added diisopropylethylamine (0.66 ml) and 4-chloro-cinnamyl chloride (467 mg). The resulting solution was stirred at room temperature for 12 hours, diluted with ethyl acetate, washed with water, brine, dried (Na₂SO₄) and concentrated
10 *in vacuo*. The residue was purified by silica gel chromatography (eluent cyclohexane:ethyl acetate 6:4) to afford 1-methanesulfonyl-5-chloro-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indoline-3,3'-pyrrolidine] (500 mg) which was characterised by its mass and NMR spectra. MS (ES+) 437/439 (M+H⁺).

Step E: A solution of 1-methanesulfonyl-5-chloro-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indoline-3,3'-pyrrolidine] obtained in Step D (256 mg) in toluene
15 (25 ml) under argon was treated with sodium bis(2-methoxyethoxy)aluminium hydride (Red-Al, 65% in toluene, 0.67 ml) and the reaction mixture was stirred at 100°C for 1 hour, cooled to room temperature, quenched by addition of ethyl acetate (10 ml), stirred for 15 min and concentrated *in vacuo*. The residue was purified by silica gel chromatography (eluent ethyl
20 acetate:methanol 95:5) to afford 5-chloro-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indoline-3,3'-pyrrolidine] (157 mg) as a yellow oil, which was characterised by its mass and NMR spectra.

Step F: A solution of 5-chloro-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indoline-3,3'-pyrrolidine] (150 mg) and triethylamine (0.24 ml) in dichloromethane (5 ml) at 0°C under
25 argon was treated with 2-chloro-isonicotinoyl chloride (147 mg). The resulting solution was stirred at room temperature for 1 hour, poured into saturated aqueous sodium bicarbonate, extracted with ethyl acetate, dried (Na₂SO₄) and concentrated *in vacuo*. Silica gel chromatography of the residue (eluent cyclohexane:ethyl acetate 6:4) afforded the title compound, 1-(2-chloropyridin-4-yl)carbonyl-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indoline-3,3'-pyrrolidine] (95 mg), which was characterised by its
30 mass and NMR spectra. MS (ES+) 498/500 (M+H⁺).

Compound III.3 was prepared according to procedures analogous to those described in Example 3.

EXAMPLE 4

5 This Example illustrates the pesticidal/insecticidal properties of compounds of formula (I).

Test against were performed as follows:

Spodoptera littoralis (Egyptian cotton leafworm)

Cotton leaf discs were placed on agar in a 24-well microtiter plate and sprayed with test solutions at an application rate of 200 ppm. After drying, the leaf discs were infested with 5
10 L₁ larvae. The samples were checked for mortality, repellent effect, feeding behaviour, and growth regulation 3 days after treatment (DAT). The following compounds gave at least 80% control of *Spodoptera littoralis*:

III-49.

15

Heliothis virescens (Tobacco budworm):

Eggs (0-24 h old) were placed in 24-well microtiter plate on artificial diet and treated with test solutions at an application rate of 200 ppm by pipetting. After an incubation period of 4 days, samples were checked for egg mortality, larval mortality, and growth regulation. The
20 following compounds gave at least 80% control of *Heliothis virescen*:

III-3, III.49

Plutella xylostella (Diamond back moth):

24-well microtiter plate (MTP) with artificial diet was treated with test solutions at an
25 application rate of 18.2 ppm by pipetting. After drying, the MTP's were infested with larvae (L₂)(10-15 per well). After an incubation period of 5 days, samples were checked for larval mortality, antifeedant and growth regulation. The following compounds gave at least 80% control of *Plutella xylostella*:

III-49.

30


Aedes aegypti (Yellow fever mosquito):

10-15 *Aedes* larvae (L2) together with a nutrition mixture are placed in 96-well microtiter plates. Test solutions at an application rate of 2ppm are pipetted into the wells. 2 days later, insects were checked for mortality and growth inhibition. The following compounds gave at least 80% control of *Aedes aegypti*

5 III-49.

CLAIMS

1. A method of combating and controlling insects, acarines, nematodes or molluscs which comprises applying to a pest, to a locus of a pest, or to a plant susceptible to attack by a pest an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a compound of formula I.



wherein Y is a single bond, C=O, C=S or S(O)_m where m is 0, 1 or 2; R¹ is hydrogen, optionally substituted alkyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, aminocarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted heterocyclyloxy, cyano, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, formyl, optionally substituted heterocyclyl, optionally substituted alkylthio, NO or NR¹³R¹⁴ where R¹³ and R¹⁴ are independently hydrogen, COR¹⁵, optionally substituted alkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl or R¹³ and R¹⁴ together with the N atom to which they are attached form a group -N=C(R¹⁶)-NR¹⁷R¹⁸; R¹⁵ is H, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted aryl, optionally substituted aryloxy optionally substituted heteroaryl, optionally substituted heteroaryloxy or NR¹⁹R²⁰; R¹⁷, R¹⁸ and R¹⁸ are each independently H or lower alkyl; R¹⁹ and R²⁰ are independently optionally substituted alkyl, optionally substituted aryl or optionally substituted heteroaryl; R² and R³ are independently hydrogen, halogen, cyano, optionally substituted alkyl, optionally substituted alkoxy or optionally substituted aryl;

each R^4 is independently halogen, nitro, cyano, optionally substituted C_{1-8} alkyl, optionally substituted C_{2-6} alkenyl, optionally substituted C_{2-6} alkynyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted C_{3-7} cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted alkylthio or $R^{20}R^{21}N$ where R^{20} and R^{21} are, independently, hydrogen, C_{1-8} alkyl, C_{3-7} cycloalkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-7} cycloalkyl(C_{1-4})alkyl, C_{2-6} haloalkyl, C_{1-6} alkoxy(C_{1-6})alkyl, C_{1-6} alkoxycarbonyl or R^{20} and R^{21} together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which may be optionally substituted by one or two C_{1-6} alkyl groups, or 2 adjacent groups R^4 together with the carbon atoms to which they are attached form a 4, 5, 6, or 7 membered carbocyclic or heterocyclic ring which may be optionally substituted by halogen; n is 0, 1, 2, 3 or 4;

each R_a is independently hydrogen, halogen, hydroxy, cyano, optionally substituted C_{1-8} alkyl, optionally substituted C_{2-6} alkenyl, optionally substituted C_{2-6} alkynyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted C_{3-7} cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted alkylthio, optionally substituted arylthio or $R^{22}R^{23}N$ where R^{22} and R^{23} are, independently, hydrogen, C_{1-8} alkyl, C_{3-7} cycloalkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-7} cycloalkyl(C_{1-4})alkyl, C_{2-6} haloalkyl, C_{1-6} alkoxy(C_{1-6})alkyl, C_{1-6} alkoxycarbonyl or R^{22} and R^{23} together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which may be optionally substituted by one or two C_{1-6} alkyl groups, or two R_a groups on the same carbon atom are $=O$ or two adjacent R_a groups are a bond, or two R_a groups together with the carbon to which they are bound form a three- to seven-membered ring, that may be saturated or


unsaturated, and that may contain one or two hetero atoms selected from the group consisting of N, O and S, and which may be optionally substituted by one or two C₁₋₆ alkyl groups; or two R_a groups together form a group -CH₂-, -CH=CH- or -CH₂CH₂; p is 0, 1, 2, 3, 4, 5 or 6; q is 0, 1, 2, 3, 4, 5 or 6; provided that when p is 2 then q is not 2;

p+q is 1, 2, 3, 4, 5 or 6;

R⁸ is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted aryl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl or optionally substituted alkenylcarbonyl; or salts or N-oxides thereof.

2. An insecticidal acaricidal and nematocidal composition comprising an insecticidally, acaricidally or nematocidally effective amount of a compound of formula I as defined in claim 1.

3. A compound of formula I'



wherein Y is CO, R² and R³ are both hydrogen, R¹ is hydrogen, optionally substituted alkyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, aminocarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted heterocyclyloxy, cyano, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, formyl, optionally substituted heterocyclyl, optionally

substituted alkylthio, NO or $\text{NR}^{13}\text{R}^{14}$ where R^{13} and R^{14} are independently hydrogen, COR^{15} , optionally substituted alkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl or R^{13} and R^{14} together with the N atom to which they are attached form a group $-\text{N}=\text{C}(\text{R}^{16})-\text{NR}^{17}\text{R}^{18}$; R^{15} is H, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted aryl, optionally substituted aryloxy optionally substituted heteroaryl, optionally substituted heteroaryloxy or $\text{NR}^{19}\text{R}^{20}$; R^{17} , R^{18} and R^{19} are each independently H or lower alkyl; R^{20} and R^{21} are independently optionally substituted alkyl, optionally substituted aryl or optionally substituted heteroaryl;

each R^4 is independently halogen, nitro, cyano, optionally substituted C_{1-8} alkyl, optionally substituted C_{2-6} alkenyl, optionally substituted C_{2-6} alkynyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted C_{3-7} cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted alkylthio or $\text{R}^{20}\text{R}^{21}\text{N}$ where R^{20} and R^{21} are, independently, hydrogen, C_{1-8} alkyl, C_{3-7} cycloalkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-7} cycloalkyl(C_{1-4})alkyl, C_{2-6} haloalkyl, C_{1-6} alkoxy(C_{1-6})alkyl, C_{1-6} alkoxycarbonyl or R^{20} and R^{21} together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which may be optionally substituted by one or two C_{1-6} alkyl groups, or 2 adjacent groups R^4 together with the carbon atoms to which they are attached form a 4, 5, 6, or 7 membered carbocyclic or heterocyclic ring which may be optionally substituted by halogen; n is 0, 1, 2, 3 or 4;

each R_a is independently hydrogen, halogen, hydroxy, cyano, optionally substituted C_{1-8} alkyl, optionally substituted C_{2-6} alkenyl, optionally substituted C_{2-6} alkynyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted C_{3-7} cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy,

optionally substituted alkylthio, optionally substituted arylthio or $R^{22}R^{23}N$ where R^{22} and R^{23} are, independently, hydrogen, C_{1-8} alkyl, C_{3-7} cycloalkyl, C_{3-6} alkenyl, C_{3-6} alkynyl, C_{3-7} cycloalkyl(C_{1-4})alkyl, C_{2-6} haloalkyl, C_{1-6} alkoxy(C_{1-6})alkyl, C_{1-6} alkoxycarbonyl or R^{22} and R^{23} together with the N atom to which they are attached
5 form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which may be optionally substituted by one or two C_{1-6} alkyl groups, or two R_a groups on the same carbon atom are =O or two adjacent R_a groups are a bond, or two R_a groups together with the carbon to which they are bound form a three- to seven-membered ring, that may be saturated or
10 unsaturated, and that may contain one or two hetero atoms selected from the group consisting of N, O and S, and which may be optionally substituted by one or two C_{1-6} alkyl groups; or two R_a groups together form a group $-CH_2-$, $-CH=CH-$ or $-CH_2CH_2-$; p is 0, 1, 2, 3, 4, 5 or 6; q is 0, 1, 2, 3, 4, 5 or 6; provided that when p is 2 then q is not 2;


15 $p+q$ is 1, 2, 3, 4, 5 or 6;

R^8 is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted aryl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl or optionally substituted
20 alkenylcarbonyl; or salts or N-oxides thereof provided that when n is 0, p is 1, q is 2, R^1 is CH_3 and all groups R_a are H then R^8 is not H, methyl, benzyl or $CH_2-CH=CH_2$ and when n is 0, $(CR_{a2})_p$ is CH-phenyl, $(CR_{a2})_q$ is $(CH_2)_2$ and R^1 is methyl then R^8 is not $COOCH_3$.

ABSTRACT
CHEMICAL COMPOUNDS

The use of a compound of formula I

5



wherein Y is a single bond, C=O, C=S or S(O)_m where m is 0, 1 or 2;

R¹, R², R³, R⁴, R⁸ and Ra are specified organic groups and p is 0, 1, 2, 3, 4, 5 or 6;

q is 0, 1, 2, 3, 4, 5 or 6; provided that when p is 2 then q is not 2;

10 p+q is 1, 2, 3, 4, 5 or 6; or salts or N-oxides thereof or compositions containing them and their using in controlling insects, acarines, nematodes or molluscs. Novel compounds are also provided.